27. *Ixias marianne*.

Papilio marianne, Cramer, Pap. Exot. iii. pl. ccxvii. C-E (1782).

Ixias bebryce, Hübner, Verz. bek. Schmett. p. 95 (1816).

Ixias agnivera, Moore, Ann. & Mag. Nat. Hist. ser. 4, vol. xx. p. 50 (1877).

Ixias depalpura, Butler, P. Z. S. 1883, p. 153, pl. xxiv. figs. 6, 7.

Ixias meridionalis, Swinhoe, P. Z. S. 1885, p. 140, pl. ix. fig. 5.

Ixias cumballa, Swinhoe, t. c. p. 141, pl. ix. figs. 13, 14.

We have twenty-five examples of the wet-season form (*I. cumballa*), thirteen of the intermediate form (*I. marianne*), twenty-three of the early dry form (*I. meridionalis*), and sixteen of the late dry form (*I. agnivera*=*depalpura*)—seventy-seven examples in all.

28. *Ixias nola*.

Ixias nola, Swinhoe, P. Z. S. 1889, p. 399.

Mahableshtar. Twelve examples. B. M.

The seasonal forms are all represented in our series; a pair of the wet and a pair of the dry form are equally labelled as types.

XXIII.—On a Precaudal Vertebra of *Ichthyosaurus australis*,
McCoy. By R. ETHERIDGE, Jun., Curator*.

THE subject of this paper is the imperfect vertebra of a large Ichthyopterigian, referable, I believe, to *Ichthyosaurus australis*, McCoy †. The original was brought under my notice by the Rev. M. Kirkpatrick, of Bega, N. S. Wales, who obtained it from Marathon, Central Queensland. With his permission a cast was taken for the Australian Museum collection. As Sir F. McCoy's description was very brief, an extended notice of one of the middle trunk, or anterior precaudal, vertebræ may be acceptable to Australian investigators.

The specimen is the centrum of a large vertebra measuring 5 inches in its vertical and transverse diameters, and rivals in size those of the gigantic *I. campylodon*, Carter, from the European Chalk, the vertebra figured ‡ by the late Sir Richard Owen measuring only 4 inches high. Our example is devoid of the neural spine, neurapophyses, and pleurapophyses, but

* From the 'Records of the Australian Museum,' vol. iii. no. 3 pp. 66-68.

† Trans. Roy. Soc. Vict. viii. 1868, p. 41.

‡ Owen, Mon. Foss. Reptilia Cret. Formation, p. 79, pl. xxii.

having the articular surfaces of the first and last well displayed. The positions of the diapophysial and pleurapophysial articular surfaces leads to the belief that the vertebra is one of the middle trunk series. It is subcircular in outline, slightly narrowed and contracted neurally. Measured across the articular surfaces from the neural to the hæmal margins the diameter is exactly 5 inches, and in a transverse direction from diapophysis to diapophysis it is an eighth of an inch short of a similar measurement. Longitudinally measured between the dia- and pleurapophysial tubercles the centrum is exactly 2 inches, but on the hæmal surface it is a quarter of an inch more.

The concave terminal articular surface visible is deep, terminating in a central fossa, the extent of the concavity being well exemplified by the matrix cast of the anterior cavity of the succeeding vertebræ at the posterior end of this specimen. This mass of matrix represents the "elastic capsule" that intervened between the vertebræ, and retains on its surface portions of the osseous tissue of the succeeding centrum. The periphery or immediate articular rim at each end is narrow, the surface thence sloping rapidly inwards, but the edges of the rims project slightly outwards, thus rendering the longitudinal or lateral surfaces of the centrum somewhat concave. The depth of the concavities is an inch, or perhaps a little more, and a longitudinal section of the centrum would be, in consequence, of a strongly hour-glass-shaped outline. The floor of the myelonal canal is $\frac{3}{4}$ of an inch wide, the joint-faces of the neurapophysial surfaces rather triangular on very strongly raised fore and aft synchondrosial articular elevations; the space between these and the diapophysial tubercles is roughly 3 inches, the latter having descended in close contiguity to the parapophysial tubercles. It is clear, therefore, that this vertebra, from the wide disassociation of the neural- and diapophyses, occupied a position in the column certainly more than one third of the trunk from the head, and, according to Owen's measurements, was near about the forty to forty-fifth vertebra, for in this region in *Ichthyosaurus* the dia- and parapophyses form a pair of separate tubercles on each side near the anterior end of the centrum.

The diapophyses are set further in from the anterior articular edge than the parapophyses; these are close to the latter, but are not connected with it by a "neck." Both are represented by large and strong rounded tubercles, separated from one another by an interval of two-eighths of an inch, this interstitial surface being deep and groove-like. The hæmal surface is quite plain.

The posterior concave articular surface is infilled with matrix, affording a complete cast of the next succeeding anterior cup, and even retaining a portion of the osseous tissue of the latter adhering to it. This tissue throughout the centrum is well preserved and dense.

The specimen is certainly of the *Campylodont* group of *Ichthyosauri*, and occupies an intermediate position in outline between an "early posterior dorsal" and a "late posterior dorsal" vertebra of *I. trigonus*, Owen*.

The largest *Ichthyosauri* attained a length of from 30 to 40 feet, and the present meridional species was in no way inferior to its gigantic fellows of the European Secondary seas. If we apply a similar method of arriving at the comparative size of an *Ichthyosaurus* as that adopted by Prof. Owen—that the jaw was "thirteen times the length of the vertical diameter of an abdominal or anterior caudal centrum"—we see in the present fossil the representative of an animal possessing a jaw a little over 5 feet in length—thus, $13 \times 5'' = 65'' = 5' 5''$ long. Prof. McCoy computed † the remains of his type specimen to represent an animal 25 feet long. Similar vertebræ to that now described he states possessed a diameter of 4 inches, and elsewhere he remarks ‡ that the longitudinal measurement reached $1\frac{1}{2}$ inch. The elastic capsule was also preserved in some of his specimens.

Mr. R. Lydekker, in the previously mentioned 'Catalogue,' gives a list of species that "cannot be classified"§. Amongst these are *I. australis*, McCoy, and *I. marathonensis*, mihi. I am afraid he has overlooked Sir Frederick's principal paper on his *I. australis*, wherein, although the description is meagre, the latter specially compares the teeth of his fossil to those of *I. campylodon*, and says they "have a rough bony square base like those of *I. campylodon* (Carter)." As regards *I. marathonensis*, mihi, less can perhaps be definitely said; but the whole of its structure, so far as we know it, is also after the type of *I. campylodon*. In my paper on this fossil I called attention to the necessity of affording another name to *I. australis*, Hector, a New Zealand species distinct from McCoy's. This has now been done by Mr. Lydekker terming it *I. Hectori* ||; but unfortunately the species is of no value, from the absence of either description or figure, all

* Lydekker, Brit. Mus. Cat. Foss. Reptilia and Amphibia, 1889, pt. 2, p. 26, figs. 13 and 14.

† Trans. Roy. Soc. Vict. ix. 2, 1869, p. 77.

‡ Ann. & Mag. Nat. Hist. (3) xix. 1867, p. 355.

§ Loc. cit. p. 113.

|| Loc. cit. p. 113.

that Sir James Hector says about it being "this genus is only represented in the collection by a single vertebral centrum."

Ichthyosaurus indicus, Lyd.*, seems to be an allied species to *I. australis*, and also vied with *I. campylodon* in size. It is from the Ootatoor Group, the homotaxial equivalent of the Chalk Marl and Upper Greensand of England.

McCoy's original specimens were from Walker's Table Mountain on the Flinders River. The present vertebra is, as before said, from Marathon on the same stream. Both are localities in the Rolling Downs Formation, or Lower Cretaceous.

XXIV.—*Description of some new Species of Heterocera.*
By HERBERT DRUCE, F.L.S. &c.

Fam. *Syntomidæ*.

Syntomis cærulescens, sp. n.

Female.—Primaries black, with six hyaline white spots—the first close to the base, the second at the end of the cell, the third below the second slightly nearer the base, the other three spots forming a band nearer the apex, the middle spot being the smallest: secondaries black, with two hyaline white spots, the first close to the base and extending to the inner margin, the second smaller and nearer the apex. The fringe of the primaries black, with a small white patch at the apex and near the anal angle. Antennæ black, with the tips white; the head, thorax, and abdomen bluish black; the front of the head white; the tegulæ black, with a small white dot at the base; abdomen banded with white at the base; the legs bluish black.

Expanse $1\frac{1}{4}$ inch.

Hab. East Africa, Delagoa Bay (*Monteiro, Mus. Druce*).
Two specimens.

Syntomis brithyris, sp. n.

Female.—Primaries smoky black, with two hyaline spots along the costal margin, the first at the end of the cell, the second beyond near the apex, below which is a large pale yellow spot; a small hyaline streak near the base, on the lower side of which is a rather wide curved yellow band that extends from the base almost to the anal angle: secondaries

* Pal. Indica (4) i. 3, 1879, p. 27.

Etheridge, Robert. 1898. "XXIII.—On a precaudal vertebra of *Ichthyosaurus australis*, McCoy." *The Annals and magazine of natural history; zoology, botany, and geology* 1, 143–146. <https://doi.org/10.1080/00222939808677943>.

View This Item Online: <https://www.biodiversitylibrary.org/item/63770>

DOI: <https://doi.org/10.1080/00222939808677943>

Permalink: <https://www.biodiversitylibrary.org/partpdf/60048>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.