Palæarctic Lepidoptera which Mr. South has catalogued in the volume before us. A short preface gives an account of Mr. Leech's life, travels, and collections; and from it we learn that he successively visited Para, Marocco, the Canaries, Madeira, China, Corea, Japan, and the North-western Himalayas, everywhere collecting and observing.

The collection, as presented by Mrs. E. Leech to the British Museum and catalogued by Mr. South, consists of 18,000 specimens, representing 1100 species. The most valuable portion consists of the materials used in the preparation of Mr. Leech's great work on the butterflies of China, Japan, and the Corea, and of the fine series of varieties and aberrations, chiefly European, a selection of the latter being represented on the two coloured plates of the volume.

Those who knew Mr. Leech personally will be pleased with the excellent portrait which forms the frontispiece; and it is a matter for real thankfulness to entomologists that Mr. Leech was not only enabled to do such excellent scientific work during his lifetime, but that his valuable collection should have found a permanent resting-place in the Natural History Museum at South Kensington after his death.

MISCELLANEOUS.

On the Evolution of the Proboscidea. By C. W. ANDREWS, D.Sc.

UNTIL the author's recent discoveries of primitive Proboscidea in the Middle and Upper Eocene formations of the Fayum, Egypt, the oldest known members of this mammalian order were *Dinotherium Cuvieri* and *Tetrabelodon angustidens*, from the base of the Miocene in France. The new Egyptian fossils not only reveal for the first time the early history of the order, but also provide more satisfactory material for the discussion of its evolution than has hitherto been available.

The most important changes in the Proboscidea occur in the skull, mandible, and dentition.

Owing to the increase in the size of the tusks and to the presence of the proboscis, the facial region of the skull becomes shortened, and at the same time the premaxillæ become wider. The presence of the proboscis also accounts for the position of the external nares. The demand for a greater surface of attachment for the muscles supporting a skull rendered heavy by the tusks and trunk is met by the great development of the diploë in certain of the cranial bones, resulting in the enormous expansion of the forwardly sloping occipital surface. The maxillæ become greatly enlarged concomitantly with the increase in the size and degree of hypselodonty of the molars. At the same time the zygomatic arch becomes weaker and the jugal takes a smaller share in its composition.

The mandible is at first short and stout, with a massive symphysis. Afterwards it becomes more and more elongated as the stature of the animals increases; and this elongation is for the most part effected by the lengthening of the symphysial region, though the backward rotation of the ascending ramus tends to the same end. The prolongation of the mandible beyond the premaxillæ must have been covered by a proboscis-like structure composed of the upper lip and nose, probably more or less prehensile at its extremity. The lengthening of the mandible seems to have reached its maximum degree in the Middle Miocene, after which it again became shortened by the reduction of the symphysis, while the fleshy and now mobile proboscis was left behind as the sole organ of prehension.

In the upper dentition the chief changes are the loss of incisors nos. 1 and 3 and the great increase in size of incisor no. 2, which eventually forms the great tusk characteristic of the later Proboscidea. The canines are soon lost. In the earliest forms some at least of the cheek-teeth (milk-molars) are replaced by premolars in the usual manner, and these teeth remain in wear simultaneously with the true molars; but in later forms no vertical succession takes place, and as the milk-molars are worn they are shed, being replaced from behind by the forward movement of the molars. Of these also the anterior may be shed, until at length in old individuals of the later types the last molar is alone functional. The gradual increase in the complexity of the Proboscidean molars is one of their most striking characteristics. All stages can be traced between the simple, brachyodont, bilophodont (quadritubercular) molars of *Moeritherium* (Middle Eocene) to the extraordinarily complex type of tooth found in *Elephas*. Thus in *Palæomastodon* (Upper Eocene) the molars are trilophodont, and the same is true of the first and second molars of *Tetrabelodon* (Miocene), in which, however, the last molar is complicated by the addition of further transverse crests. In the Stegodonts of the Siwalik Hills (Pliocene) a further increase in the number and height of the crests takes place, and the whole crown of the tooth is more or less covered with a thick coat of cement. Still later the transverse crests become highly compressed laminæ united by cement, and these are as many as twenty-seven in number in the Pleistocene *Elephas primigenius* and the recent *E. indicus*.

The evolution of the lower molars corresponds with that of the upper molars. Of the lower incisors the middle and outer pairs (nos. 1 and 3) are soon lost, but the second pair remains functional for a long geological period. When the symphysis becomes shortened these incisors are sometimes retained as vestiges (*e. g.* in *Mastodon americanus*), but in the genus *Elephas* they have completely disappeared.—*Abstract of paper read before the Royal Society on March 26th, 1903.*

Andrews, Charles William. 1903. "On the evolution of Proboscidea." *The Annals and magazine of natural history; zoology, botany, and geology* 11, 527–528.
<https://doi.org/10.1080/00222930308678813>.

View This Item Online: <https://www.biodiversitylibrary.org/item/63777>

DOI: <https://doi.org/10.1080/00222930308678813>

Permalink: <https://www.biodiversitylibrary.org/partpdf/60214>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.