jection of the indigestible portion of its food is effected by dilating the whole body with water and then ejecting it.

At the same time a small species of *Clio*, several *Beroë*, one specimen of a free *Actinia*, and a very beautiful Crustacean, forming a new genus in the family *Pontia* and allied to *Irenæus*, were obtained. The last is characterized by its large size, "the enormous length of the four central tail-filaments; the inner of which are not themselves armed with filaments, all the others being so; each of the antennæ has a joint at the distal part of the first third, thus enabling the animal to bend them and conceal them under its body."

On the second fine day a most interesting Ciliograde was obtained of very peculiar form, and having the ciliated ribs transverse instead of longitudinal; the cilia arising from either edge of the ribs. Mr. Goodsir thinks that its minute structure, which is very complicated, proves its affinity to the *Diphydæ* and also the *Salpæ*. All the *Medusæ* obtained on these two days were Ciliogrades.

On the 23rd of June, having rounded Cape Farewell in a heavy gale of wind on the preceding day, on entering Davis' Straits numerous beautiful *Pteropoda* of the genus *Clio* were obtained in company with *Spiratella*. They were swimming actively in the water and were adorned with the brightest colours; only approaching the surface of the sea on calm evenings. The observations made upon these animals have enabled Mr. Goodsir to confirm the observations of Eschscholtz in most points.

On the 25th numerous *Medusæ* were seen, all of these of the family *Beroidæ* and mostly of the species *B. punctata* of Eschscholtz. On the 27th soundings were obtained in 41 fathoms and a dredge put down, which produced, amongst many other interesting captures, a new species of *Caprella*, *Amphipoda* in great number, several *Asteriadæ*, a *Terebratula* and several other Mollusca, an Isopod forming a new genus allied to *Munna*, a very beautiful Ascidian, four species of fish—*Cyclopterus*, *Liparis*, *Ammodytes*, and a beautiful species new to the observer. On the same day a large shoal of the Caing whale (*Phocæna Melas*) passed on their way towards the south.

On the 28th a dredge was sunk to the enormous depth of 300 fathoms, and produced many highly interesting species of Mollusca, Crustacea, *Asteriadæ*, *Spatangi* and Corallines; such as *Fusus*, *Turritella*, *Venus*, *Dentalium*, &c.; and also some large forms of *Isopoda*. As bearing upon the geographical distribution of species, Mr. Goodsir considers the occurrence of *Brissus lyrifer* (Forbes) and *Alauna rostrata* (Goodsir) as of the greatest interest, both of them being natives of the Scottish seas. The remarkable depth also appears to us to give peculiar interest to

these researches, as we believe that the deepest dredgings ever previously obtained were those of Prof. E. Forbes in the Levant, the deepest of which was 230 fathoms, itself far beyond any made by other naturalists. Such valuable and laborious researches can only be made when the officers of a ship have such kindly dispositions towards the observer of nature as was the case during Prof. Forbes's voyage, and is now shown by those under the command of Sir J. Franklin.

On the 1st of July two specimens of a small species of *Beroë* were obtained, which greatly illustrated the process of their development. "A thick germinal membrane of a red colour was observed lining the central cavity of the body, in which both male and female cells appeared to be developed. The ova having arrived at some size project so far as to become pedunculated, and so hang from the membrane into the cavity. The male cells are also developed in the same membrane."

Mr. Goodsir is making minute observations upon the ice of the bergs, and as he purposes continuing them throughout the voyage, there can be little doubt of his arriving at valuable conclusions. He observes that it is quite without salt; this was to be expected, when we consider that they are not formed of marine ice, but are parts of glaciers which have become detached and fallen into the sea. "The surface of a mass when melting presents numerous flat concavities, all of them of about the same size and form, and without any interruption, excepting the ridges forming the walls of separation." A loud cracking noise is heard and small particles are occasionally thrown off. The minute structure of the ice consists of three series of cells—two traversing the mass in one direction, and one at right angles to them. Of the former, one series consists of moderately large and quite globular cells of nearly uniform size, each having within it "a small globule of a peculiar fluid;" the oblong, sausage-shaped cells of the other series also contain small globules, but usually several instead of one. The mode of formation of these cells and the nature of the inclosed fluid are subjects to which his attention will be especially turned. The third series consists of very minute cells, arranged in well-defined wavy bands, which run across the lines formed by the other series. These bands are of an opake white colour.

We also find some observations upon the action of floating ice upon the granitic shores of the islands. All the rocks below high water mark and some considerably above it are rounded off into long irregular ridges with intervening hollows by the half-floating masses of ice.

As the ships were to stay at the Whale-fish Islands for a few

Thomas, Oldfield. 1895. "XXIII.—On a new species of Eliurus." *The Annals and magazine of natural history; zoology, botany, and geology* 16, 164–165.

<https://doi.org/10.1080/00222939508680247>.

View This Item Online: <https://www.biodiversitylibrary.org/item/71834>

DOI: <https://doi.org/10.1080/00222939508680247>

Permalink: <https://www.biodiversitylibrary.org/partpdf/60350>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.