

Nondum quidem illam undique satis examinavimus, ut de illius vi atque pretio certi aliquid audeamus prædicare; verisimile vero nobis visum est, adminiculum haud minimi ponderis hanc notam præbere ad species affines dignoscendas." The application of iodine also manifests a long-desired distinction between *Sphærophoron coralloides* and *fragilis*, tinging the stratum medullare of the thallus of the former of a violet or intense blue, and that of the latter of a yellow colour.

LIV.—*On a new Species of Victorian Honey-eater.* By FREDERICK M'COY, Prof. Nat. Sc. Melbourne University, and Director of the National Museum, Victoria.

*Ptilotis Leadbeteri* (M'Coy).

The Subcrested Honey-eater.

Feathers of the crown of the head suberect, so as to form a slightly marked subcompressed crest. Bill strong, moderately arched. Colours (male):—Crown of head and nape dull greenish yellow; back, wings, and tail dull greenish black, the outer edge of the wing- and tail-feathers narrowly edged with dull greenish yellow; lores, or wide space from the gape over and under the eye, the ear-coverts, and extending to the shoulder, intense glossy black; the ear-plume large, and of the most intense pure golden yellow; cheeks below the black bright yellow; chin dull black; breast and abdomen and under tail-coverts dull yellow tinged with olive, particularly at the sides; under wing-coverts blackish, with a narrow bright-yellow edge near the shoulder; the middle of the inner edge of the primaries margined with dull whitish; all the outer tail-feathers tipped with yellowish white, chiefly on the inner circles. Bill and feet brownish black.

Total length 9 inches; bill from gape  $\frac{11}{16}$  inch, from forehead  $\frac{9}{16}$  inch; wing  $4\frac{1}{4}$  inches, tail  $4\frac{3}{8}$  inches, tarsi 1 inch.

Female: total length  $7\frac{1}{2}$  inches; with the back, wings, and tail of a dark brownish olive, where the male is black.

This splendid new Honey-eater is most allied to the *P. auricornis*, from which its greater size, subcrested head, blackish upper colouring, shorter and thicker bill, and the larger and more intensely coloured ear-plumes, relieved on the intense black extending to the shoulder, easily distinguish it. The subcrestate head, and the female differing in colour from the male, suggest a new subgeneric section for this fine bird, which I have great pleasure in naming after my able and zealous taxidermist at the Melbourne National Museum, whose great ability and diligence well deserve the compliment.





McCoy, Frederick. 1867. "LIV.—On a new species of Victorian honey-eater." *The Annals and magazine of natural history; zoology, botany, and geology* 20, 442–442.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/72234>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/61288>

**Holding Institution**

University of Toronto - Gerstein Science Information Centre

**Sponsored by**

University of Toronto

**Copyright & Reuse**

Copyright Status: NOT\_IN\_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.