like down; the peduncle is formed of spirally twisted threads, and divides below into a few rootlets:" and he believes that the sponge grows sticking in the mud.

Dr. O. Schmidt kindly sent me a slide with specimens of the spicules of *Tetilla*, but I do not find any trifurcated spicules on it; one of them is figured across the base of the sponge, t. 5. f. 10. It also belongs to *Tethyadæ*.

On *Hyalonema*, Gray.

Professor E. Perceval Wright of Dublin has just returned from Setuval, where, with the kind assistance of Prof. Bocage of Lisbon, he has succeeded in dredging living specimens of this strange organism. The *Hyalonema*-ground is in a valley, some thirty miles to sea, south-west of Setuval, and is from 400 to 500 fathoms in depth. Prof. Wright "regards the siliceous axis as the stem of the "sponge-mass called *Carteria* by Dr. J. E. Gray, and has determined that the end of the axis, where the fibres become loose, is "that one imbedded in the mud, the sponge-mass being on the "summit, and presenting forms of very various outline. The "sponge-mass is provided with a number of oscula looking upwards, "these being covered over by a beautiful open network of spicules. "When the sponge-mass is washed away or destroyed, the parasitic "*Palythoa*, which was seen living, and in the act of protruding its "tentacles, grows up over that portion of the siliceous axis which is "left uncovered by the mud; but numerous examples of the siliceous stem exist uncovered by the parasite. The Lisbon Museum "has now, thanks to Prof. Bocage, the most magnificent series of "this sponge in the world." Prof. Wright will shortly publish fuller details of this interesting discovery.

Castle, Dublin, Sept. 22nd.

MY DEAR DR. GRAY,—Many thanks for your kind note, which I got on my return from the expedition Carpenter and I made to the North Sea * * * * *

Now, as to our expedition. In the mud of the Gulf Stream (at 550 fathoms) we got *Hyalonema* living *upside down*, as I already suspected from Lovén's paper; but, besides *Hyalonema*, we got at least half a dozen new forms of vitreous sponges, most remarkable, and some of them as beautiful as the flower-basket.

Of these you will, of course, get specimens; but in the first place I must clean and prepare them and describe them for the 'Phil. Trans.'

In another locality we got *Brisinga* and the wonderful little Crinoid *Rhizocrinus* * * * * *

Ever truly yours,
WYVILLE THOMSON.

1868. "On Hyalonema, Gray." *The Annals and magazine of natural history; zoology, botany, and geology* 2, 320–320.

<https://doi.org/10.1080/00222936808695811>.

View This Item Online: <https://www.biodiversitylibrary.org/item/72302>

DOI: <https://doi.org/10.1080/00222936808695811>

Permalink: <https://www.biodiversitylibrary.org/partpdf/61343>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.