"283. LAPLYSIA. *Corpus repens, obvelatum membranis reflexis.*

"*Clypeo dorsali, membranaceo, pulmones obtegente.*

"*Foramen laterale, dextrum, pro genitalibus.*

"*Anus supra extremitatem dorsi.*

"*Tentacula quatuor, anterieus sita.*

"*depilans.* 1. LAPLYSIA.

"*Syst. Nat.* 10. p. 653. *Tethys limacina.*

"*Rond. pisc.* 1. p. 520. *Lepus marinus.*

"*Gesn. aquat.* 475. *Lepus marinus Rondeletii.*

"*Bohads. mar.* 3. t. 1, 2, 3. *Lerneæ graphice.*

"*Seb. mus.* 3. t. 1, f. 8, 9.

"*Habitat in M. Mediterraneo; sanie depilans tactu.*

"(B. 51.) *foetidissima ad nauseam usque.*"

The description of the genus is implied for the species *depilans*, and it is also said to be the *Tethys limacina* of the tenth edition *. The second reference is to the same figure of Rondelet formerly cited for *Tethys leporina*. The third reference repeats the earlier citation to Gesner, with the same mistake as to the page. The fourth reference is to the excellent figure of Bohadsch's *Lerneæ*, representing unmistakably the *Aplysia depilans* of authors. The reference to Seba is less happy, the figures being too ambiguous for certain determination. It is perfectly evident that Linnæus's generic characters of *Laplysia* were derived from Bohadsch's work; and as the best figures were from the same source, the traditional identification of *depilans* is fully sustained.

SUMMARY.—From the foregoing facts it would appear that (1) the generic name *Tethys*, Linn. 1758, must replace *Aplysia* and *Laplysia*, Linn. 1767; and (2) as a substitute for *Tethys*, Linn. 1767, not 1758, we will probably be compelled to adopt either one of the new spellings of this name proposed in the early part of the century or an entirely new generic term.—*Proc. Acad. Nat. Sci. Philad.* August 27, 1895, pp. 347-350.

On the Origin of the Triradiate Spicules of Leucosolenia.

By E. A. MINCHIN.

In *Leucosolenia coriacea* the youngest spicules are found to be surrounded by six cells, which are similar in all their characters to the cells of the external flat epithelium of the sponge, and undoubtedly derived from this layer. It appears that three cells of the external epithelium wander inwards, and give rise to six by division of each cell into two, the six cells being arranged in such a way that three are placed more internally, *i. e.* towards the gastral

sight, for the first use of the word, on page 1072 of the *Syst. Nat.* 12, is in the correct form "*Aplysia*." The generic diagnosis given on this page is brief, but sufficient:—"283. *Aplysia* Tentacula 4. Anus supra postica."

* We would not replace the specific name *depilans* by *limacina*, because the latter was not recognizably defined in Linnæus's earlier edition.

surface of the body-wall, and three more externally, towards the dermal surface. Each of these sets of three cells has a form which might be compared to a trefoil, and the whole mass may be described as two such trefoils superposed, the cells of one trefoil exactly corresponding to those of the other.

The spicule is formed by the three inner cells, a ray being formed by each cell. In many instances it appears as if the three rays were formed quite separately and afterwards fused at the centre.

The three outer cells soon lose their rounded form, and, by throwing out processes, assume an amœboid appearance. After the spicule-rays have attained a length of 10 or 15 μ , the three outer cells are no more to be found, having apparently rejoined the flat epithelium from whence they came. The three inner cells alone secrete the rays, and continue to do so until the spicule is full-grown.

The spicule-rays soon appear to project beyond their formative cells, but are in reality covered by a thin layer of protoplasm. At the same time the spicule-sheath makes its appearance as a denser layer of substance between the protoplasm of the formative cell and the calcareous spicule, and it is by continued calcification of the sheath that the spicule grows.

The spicule-rays attain their full thickness at their bases before they have reached their full length. The formative cells remain at the bases of the rays until this portion is built up to its full thickness. Each formative cell then migrates along its ray towards the tip, building up the ray to its full thickness as it goes. In the fully formed ray the formative cell is found adherent to the extreme tip.

Theoretical considerations.

(1) The origin of the spicule-forming cells—that is to say, of the whole connective-tissue system in these sponges—from the external flat epithelium, is another nail in the coffin of the so-called mesoderm in these forms. Sponges are to be regarded as two-layered animals, composed of a dermal and a gastral layer. The dermal layer is differentiated into (1) an external flat contractile epithelium, the neuro-muscular system, and (2) an internal connective-tissue layer. The gastral layer consists of the collar-cells. The amœboid wandering cells are perhaps also to be reckoned with the gastral layer.

(2) The fact that each ray of a triradiate spicule is formed by a single cell shows that each triradiate spicule must be regarded as derived from the fusion of three originally separate monactinal spicules. This supports Schulze's theory, namely, that the tri-radiates of the more primitive Ascons have arisen as an adaptation to the structure of the sponge, and goes against Dreyer's theory that the primitive spicule of all sponges is a tetraxon, a form explained by him as the direct mechanical outcome of the vesicular structure of living bodies.—*Proc. Roy. Soc.* vol. lviii. no. 350, pp. 204, 205.

THE ANNALS

AND

MAGAZINE OF NATURAL HISTORY.

[SIXTH SERIES.]

No. 96. DECEMBER 1895.

LVIII.—*The Pectoral and Pelvic Girdles of Murænosaurus plicatus.* By C. W. ANDREWS, F.G.S., Assistant in the British Museum (Natural History).

PHILLIPS, in the 'Geology of Oxford' (p. 313), described under the name *Plesiosaurus plicatus* certain cervical and dorsal vertebræ from the Oxford Clay of Shotover. From the same horizon in the neighbourhood of Peterborough numerous more or less complete skeletons of a species of Plesiosaur have been collected in which the vertebræ are of a similar character. These specimens present a considerable range of variation both in size and in the degree to which fusion of the elements of the vertebræ has taken place; in some the neural arches and cervical ribs are still quite free, in others they are co-ossified with the centra. It was on one of the skeletons in which the latter condition obtains that Professor Seeley founded his description of *Murænosaurus Leedsi*. The genus *Murænosaurus* is distinguished from *Cryptoclidus*, to which the greater number of specimens from this locality are referable, by the enormous length of the neck (about seven times that of the head), caused not only by the much larger number (forty-four) of cervical vertebræ, but also by the greater length of the individual centra. There are also numerous differences in the pectoral girdle and limb-bones which fully establish the right of this type of Plesiosaurian skeleton to generic rank.

With regard to the species there is some difficulty. Mr. Lydekker has included under *Cimoliosaurus plicatus* Phillips's species *Plesiosaurus plicatus*, *P. hexagonalis*, and *P. infraplanus*, as well as Professor Seeley's *Murænosaurus Leedsi*. For the specimen (Leeds Coll. 28), the limb-girdles of which are figured in the present paper, he has suggested the specific name *Cimoliosaurus durobrivensis*, considering it separable from *C. plicatus* on account of its somewhat shorter cervical vertebræ. Comparison of some six or seven more or less complete sets of cervical vertebræ (including those of the type specimens of *Murænosaurus Leedsi* and *Cimoliosaurus durobrivensis*) with one another shows that, apart from variations in size and in the extent to which the arches and ribs have fused with the centra, no important differences are to be found. The size and condition of ossification are probably dependent merely on the age and sex of the individual. It is, however, possible that the comparatively small individuals in which fusion between the arches and centra has already occurred may be specifically distinct, in which case the name *Murænosaurus Leedsi* must be applied to them, the type specimen of that species being of this form. On the other hand, as already remarked, these differences may be merely sexual, and, as Mr. Boulenger has shown in the case of the *Chelonia* *, the fusion of other elements with the vertebral centra takes place very irregularly, and consequently is of little systematic value. For the present, therefore, it seems best to regard all the specimens possessing cervical vertebræ of the type of *Plesiosaurus plicatus* as belonging to that species.

In the present note specimens of the pectoral and pelvic girdles which have lately been mounted for exhibition at the Natural History Museum are described. These belong to the same individual (R. 2428, Leeds Coll. 28), which is that for which Mr. Lydekker proposed the name *Cimoliosaurus durobrivensis*.

The Pectoral Girdle. (Fig. 1.)

In general form the *scapula* (*sc*) resembles that of *Cryptoclidus*, but differs from it in several points, *e. g.* (1) the posterior border of the dorsal ramus near its upper end bears a sharp crest, apparently for the insertion of muscle; (2) the dorsal ramus widens out much more rapidly towards its base and its anterior edge is thin and sharp; (3) the ventral ramus is considerably wider, and also has a comparatively thin anterior

* Trans. Zool. Soc. vol. xiii. pt. 8, p. 309.

edge, which bears no tubercle such as occurs in *Cryptoclidus* (see this Magazine, ser. 6, vol. xv. p. 337, fig. 2 A, t). The median portion of the ventral bars had already begun to grow back to form a junction with the anterior prolongations of the coracoids, but were still separated from them by a considerable interval; other pectoral girdles in the collection show this junction. The scapulæ had not yet quite met in median symphysis, and anteriorly were separated from one another by a wide V-shaped interval. In life this space was wholly or partially closed by the clavicular arch, which unfortunately is wanting in this specimen. The scapulæ anteriorly curve inwards and upwards in such a way that, with the clavicles, they formed the anterior wall of the thorax.

The general form of the *coracoids* (*cor*) is shown in the figure (fig. 1). Between the glenoid surfaces the bones are very thick and massive, and unite in a strong median symphysis

Fig. 1.

Pectoral girdle of *Muraenosaurus* from above. *sc*, scapula; *cor*, coracoid; *h*, humerus. About $\frac{1}{10}$ nat. size.

about 5 centim. deep at the thickest part. In this region the visceral surface of the coracoids is concave from side to side, forming an arc of a circle whose radius is about 28 centim. Behind this strong transverse ridge the visceral surface is depressed, owing to the rapid thinning of the bone, which, except near the outer border, is less than a centimetre thick. In this thin posterior region the coracoids meet at an angle of about 120° . The outer concave border is somewhat thickened, but is not produced backwards and outwards into

projecting processes as in the fully ossified coracoid of *Cryptoclidus*. The thin inner posterior portion of the bone is frequently broken away, and the remaining portion, consisting of the thickened anterior and external borders, has then the form shown by Prof. Seeley in his figure of the coracoids of *Murænosaurus Manselli* *.

The posterior convex margin in the present specimen bears rough grooved surface for cartilage, showing that growth was still in progress in this direction. Anteriorly the coracoids are prolonged forwards in the middle line in short pointed processes, which extend towards the backward processes of the scapulæ, but do not reach them; both bones, however, bear surfaces for cartilage, and were probably already united by a bar of that substance.

The dimensions of the pectoral girdle are :—

	centim.
Extreme length from anterior end of scapulæ to hinder border of coracoids	59
Width across coracoids in a straight line between the hinder edges of the glenoid facets.....	38
Extreme length of coracoids	40
Width of coracoids at narrowest point.....	27

The humerus (*h*) figured measures about 31 centim. long; the smallest circumference of its shaft is 21 centim., and the width of its expanded distal extremity is 19 centim.

The Pelvic Girdle. (Figs. 2, 3.)

The pelvis represented in figs. 2 and 3 gives a clearer idea of the structure of that portion of the Plesiosaurian skeleton than any other specimen or figure with which I am acquainted.

The *pubis* (*pu*) is a broad nearly flat plate of bone, very thin except near the median border and the articular surfaces for the ischium and femur; the posterior margin is also somewhat thickened. The width of the bone from the antero-external to the antero-internal angle is 28 centim., the length from the ischial surface to the anterior border 25·5 centim. The length of the acetabular surface is about 8 and of the ischial surface 4 centim. In the specimen figured the left pubis has been restored in plaster. The two pubes seem to have made an angle of about 120° with one another; they were still separated in the middle line by a pad of cartilage, which was widest in front and behind; whether the posterior portion of this cartilage extended back to join that lying between the

* Quart. Journ. Geol. Soc. vol. xxx. (1874) p. 437, fig. 2.

Andrews, Charles William. 1895. "LVIII.—The pectoral and pelvic girdles of *Muraenosaurus plicatus*." *The Annals and magazine of natural history; zoology, botany, and geology* 16, 429–434. <https://doi.org/10.1080/00222939508680300>.

View This Item Online: <https://www.biodiversitylibrary.org/item/81045>

DOI: <https://doi.org/10.1080/00222939508680300>

Permalink: <https://www.biodiversitylibrary.org/partpdf/62783>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.