Lepidoplaga multidentalis, sp. n.

Fore wings pale yellowish; the costa, transverse lines, and stigmata fuscous; first line close to base, dentate; second from two thirds of costa to near middle of inner margin, strongly dentate and incurved above the inner margin; a straight dark fuscous shade from costa before apex to inner margin just before anal angle also strongly and regularly dentate; fringes concolorous, with a row of minute dark brown dots at their base. Hind wings paler, with a short central fuscous line and a broad shade, continuing that on fore wings, starting below the costa and ending in hind margin at one third from anal angle. Abdomen and back of thorax pale yellow; outside of palpi and front of thorax fuscous; face fuscous yellow, edged with whitish. Underside paler; both crests of scales very glossy.

Expanse of wings 35 millim.

One male from the Khasias.

Lepidoplaga uniformis, sp. n.

Fore wings uniform pale yellow, with the subcostal region slightly deeper tinged and the costa very narrowly white; lines slightly darker, indistinct, first before one third, oblique, second at two thirds, oblique and straight towards anal angle for two thirds, then curved in to below discocellular, and again at an acute angle outwards to inner margin at two thirds; no discal spot, but the fovea at end of cell clearly indicated, that towards the base hardly visible; fringe concolorous. Hind wings rather paler, with the exterior line reproduced, beyond which the ground-colour becomes greyer. Head, thorax, and abdomen concolorous with wings. Underside very glossy, the fore wings greyish, the hind wings paler.

Expanse of wings 21 millim.

One male from the Khasias.

PACHYBOTYS, gen. nov.

Fore wings elongate; costa straight; apex rounded; hind margin curved; anal angle rounded off. Hind wings narrow, with round hind margin. Antennæ simple in both sexes; labial palpi shortly porrect, laxly scaled; maxillary palpi insignificant; tongue and ocelli present; legs thickened with hairs.

A genus of insects of stouter and thicker build than *Opsibotys*, with denser scaling and quite different pattern of

markings; the two stigmata are well expressed and large; the exterior line rectangularly produced outwards in the middle, followed by a well-developed submarginal line.

Type *P. spissalis*, Guen. (*Botys*).

Pachybotys minialis, sp. n.

Wings dull vermilion-red, with dull black markings; a spot at the base of the fore wings and the whole costa blackish; first line at one third, thick, straight and oblique, parallel to hind margin; second line before two thirds, at first parallel to hind margin, then running abruptly outwards, describing a quadrangular sinus, and ending on inner margin at two thirds as a straight oblique line in a line with the costal arm; discal mark lunular; space between the lines thickly dusted with blackish, except immediately beyond discal lunule, which remains of the ground-colour; submarginal line curved, rather interrupted; fringe paler, with a dark broad basal line. Hind wings paler red, especially along costa, with traces of a dark line at two thirds, distinct only near inner margin, and of a submarginal one, much thinner; fringe reddish, with no black basal line. Head, thorax, abdomen, and legs all vermilion-red. Underside rather duller; the fore wings blackish beyond the submarginal line.

Expanse of wings 26 millim.

Three females from Queensland.

Pachybotys plenistigmalis, sp. n.

Wings deep yellow, suffused over the basal half with brownish yellow; costa at base ferruginous; lines deeper yellow, first at one fourth, curved outwards, second at three fourths, at first straight from costa, then forming a shallow sinus of three teeth, and running in to touch the base of the reniform stigma, where it forms a dark brown spot, and descends to the inner margin; traces of a curved and sinuous submarginal line; stigmata dark ferruginous, large, the first round, the second 8-shaped, with traces of a metallic centre. Hind wings with an angulated deep yellow central line and a broadish submarginal band, both lustrous; fringes of both wings yellow. Head, thorax, and abdomen dark yellow; palpi ferruginous. Underside paler yellowish white, with the stigmata and lines showing through.

Expanse of wings 26 millim.

A few from the Khasia Hills.

[To be continued.]

LXV.—*Descriptions of Five new Species of Ennea from South Africa.* By JAMES COSMO MELVILL, M.A., F.L.S., and JOHN HENRY PONSONBY, F.Z.S.

[Plate XVIII.]

THE following species of *Ennea*, in most instances considerably smaller than those hitherto described from the South-African region, were received by Mr. Wotton from the neighbourhood of Grahamstown, and by him placed in our hands for determination.

Ennea cimolia, sp. n. (Pl. XVIII. fig. 2.)

E. testa cylindracea, minuta, recta, tenui, nitida, apud apicem compressa; anfractibus sex, ventricosis, nitidis, ad suturas paullum impressis, omnino laevibus; apertura ovata, labro incrassato, pellucido, dentibus vel plicis tribus instructo, plica parietali magna, intrante, dente labiali conspicuo, plica marginem subtus columellarem semicelata, effusa.

Long. 2.75, lat. 1.25 mm.

Hab. Grahamstown.

A very delicate transparent shining little mollusk, with whorls perfectly smooth; mouth ovate; peristome with three processes, of which the parietal plait is deep-seated, the labial tooth conspicuous, the columellar plait being effuse, large, and half-concealed.

Ennea Farquhari, sp. n. (Pl. XVIII. figs. 3-5.)

E. testa minuta, rimata, dolioliformi, tenui, nitida; anfractibus sex, ventricosis, ad suturas multum impressis, undique longitudinaliter pulchre striatis; apertura ovato-triangulari, labro incrassato, dentibus vel plicis quatuor instructo, plica parietali magna, acinaciformi, acuta, valde prominente, dente labiali bipartito, plica columellari minore, simplici.

Long. 3, lat. 1 mm.

Hab. Grahamstown.

Several specimens, two being in exceptionally fine and live condition, quite transparent, with yellow-rose hue. The species being somewhat variable, we give three representations of it. It is tun-shaped, minute, thin, shining, six-whorled; whorls ventricose, beautifully longitudinally striate, beneath triangularly ovate; outer lip thickened, mouth adorned with four processes, the parietal plait being large, scimitar-shaped,

Melvill, James Cosmo and Ponsonby, John Henry. 1895. "LXV.—Descriptions of five new species of Ennea from South Africa." *The Annals and magazine of natural history; zoology, botany, and geology* 16, 478–480.

<https://doi.org/10.1080/00222939508680307>.

View This Item Online: <https://www.biodiversitylibrary.org/item/81045>

DOI: <https://doi.org/10.1080/00222939508680307>

Permalink: <https://www.biodiversitylibrary.org/partpdf/62787>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.