

LARVAL MITES (ACARI: TROMBIDIIDAE) PARASITIC ON APHIDS IN IRAN: KEY, A NEW SPECIES AND NEW RECORD

Zhang, Z.-Q. & N. Rastegari, 1996. Larval mites (Acari: Trombidiidae) parasitic on aphids in Iran: key, a new species and new record. – Tijdschrift voor Entomologie 139: 91-96, figs. 1-6. [ISSN 0040-7496]. Published 15 October 1996.

A key to larvae of Trombidiidae (Acari: Prostigmata) found ectoparasitic on aphids in Iran is presented. *Allothrombium shirazicum* Zhang sp. n. is described and illustrated from larvae parasitic on *Forda marginata* Koch (Pemphigidae) and unidentified aphids in Shiraz, Iran. *Monotrombium simplicium* Zhang is newly recorded from larvae parasitic on aphids in wheat fields in Shiraz, Iran.

Z.-Q. Zhang, International Institute of Entomology, 56 Queen's Gate, London SW7 5JR, UK.

Key words. – Acari; Trombidiidae; classification; larvae; key; ectoparasites; aphids; Pemphigidae; *Monotrombium*; *Allothrombium*; Iran.

Larvae of the genus *Allothrombium* and *Podothrombium* are common ectoparasites of aphids and are expected to have potential for use as biocontrol agents against aphids (Eickwort 1983, Welbourn 1983, Zhang 1991a, Zhang & Xin 1992). Discovery and accurate description of these parasitic mites are prerequisites for any research toward their potential use in pest control programs (Eickwort 1983, Welbourn 1983).

Recently, H. Norbakhsh of Shahid Chamran University, Ahwaz, Iran sent to the senior author some larval trombidiid mites which were found ectoparasitic on various wheat aphids in Shiraz, Iran. A study of these mites revealed a new species (*Allothrombium shirazicum* Zhang sp. n.) and a new record (*Monotrombium simplicium* Zhang) from Shiraz, Iran. The purpose of this paper is to describe the new species and to present a key to larvae of Trombidiidae parasitic on aphids in Iran.

The terminology and abbreviations used in this paper are adapted from Robaux (1974) and Welbourn & Young (1988). All the measurements of length are in micrometers.

Key to larvae of Trombidiidae parasitic on aphids in Iran

1. Coxa II with one seta; genu II and genu III each with one solenidion *Monotrombium* (*M. simplicium* Zhang)
- Coxa II with two setae; genu II and III each with two solenidia (*Allothrombium*) 2


2. Tarsus III with two normal claws and an empodium *A. pulvinum* Ewing
- Tarsus III with one normal claw and empodium; inner claw vestigial 3
3. Idiosoma with more than 20 dorsal setae and more than 10 ventral setae *A. mossi* Zhang
- Idiosoma with 20 dorsal setae and less than 10 ventral setae 4
4. Legs short; tarsi I-III < 60 µ ... *A. triticism* Zang
- Legs long; tarsi I-III > 90 µ *A. shirazicum* Zhang sp. n.

Allothrombium shirazicum Zhang sp. n.
(figs. 1-6)

Type material. – Holotype larva (ZQZ 96-0128-3a) parasitic on an aphid, collected by N. Rastagari (No. 14, 58-19), on 20.iv.1992 in Shiraz, IRAN. – Paratype larva (ZQZ 93-0128-3b), same data as holotype. Paratype larvae (ZQZ 930128-2c, d) parasitic on *Forda marginata* Koch (Pemphigidae), collected by N. R. Nowband (No. 1-19), on 14.vi.1992 in Shiraz, IRAN. All types deposited in the The Natural History Museum (BMNH), London.

Description

Larvae with the following features: fD = 4(+2)-6-4-4-2 = 18(+2); fV = 2-2-2u-2 = 8; fnTr = 1-1-1; fnFe = 5-4-4; fnGe = 4-3-3; fnTi = 5-5-5; fnTa = 17-14-13; fSol = I(0-2-2-1), II(0-2-2-1), III(0-2-0-0); fκ = I(1-1), II(1-0), III(0-0); fζ = 2-0-0; fε = 1-1-0; fPp = 0-0-0-BNN2-BBNNNω; IP = 1298 (1280-1320).



Figs. 1-3. *Allothrombium shirazicum* Zhang sp. n. Holotype larva. – 1, Idiosoma, dorsal view; 2, Idiosoma, ventral view; 3, Gnathosoma.

Larva. – Measurements are means of four specimens, with range in parentheses. Idiosoma engorged, holotype 750 long, 450 wide. Idiosoma dorsally with a scutum, a scutellum, a pair of ocular sclerites, and 24-26 dorsal setae. Scutum pentagonal in shape, widest at postero-lateral angles, with convex posterior side; small punctation on scutum denser on posterior

part than on anterior part; AM setae barbed, near antero-lateral angles of scutum; PL setae barbed, at postero-lateral angles; AL setae barbed, between AM and PL setae; S barbed only in distal half, between AL and PL setae. Scutellum with two barbed setae; wider than long, with small punctation throughout. Standard measurements of scutum and scutellum as

follows: AM 39 (29-45); AA 60 (53-66); AL 44 (43-47); AW 87 (78-90); MA 42 (41-44); PL 72 (70-73); PW 104 (100-114); AP 46 (38-53); S 84 (79-90); SB 68 (64-70); ASB 94 (90-99); PSB 56 (55-59); SD 151 (145-158); W 124 (119-130); HS 37 (33-38); LSS 86 (79-93); SL 68 (65-72); SS 36 (29-43). Ocular sclerite, 37 (34-40) long, lateral to postero-lateral angles of scutum; with 2 eyes, the anterior eye (13-15 in diameter) larger than the posterior one (9-11 in diameter). All dorsal setae barbed, arising from small setal sclerites; dorsal setal formula $fD = 4(+2)-6-4-4-2 = 18(+2)$.

Idiosoma ventrally with three pairs of coxae, 1 pair of intercoxal setae, eight pairs of ventral setae, and an anus. All setae on ventral idiosoma with barbs. Coxa I 82 (75-88) long, with two barbed setae. Coxa II 84 (81-88) long, with two barbed setae. Coxa III 74 (69-75) long, with a single barbed seta. Intercoxal setae between coxa III. Ventral setae with small setal sclerites; ventral setal formula $fV = 2-2-2u-2 = 8$.

Gnathosoma truncate posteriorly. Palpal setal formula $fPp = 0-0-0-BNN2-BBNN\omega$. A pair of adoral setae nude, 8 (6-9) long. A pair of subcapitular setae thick, branched distally, 11 (11-12) long, 119 (18-20) apart at base. Cheliceral base 65 (62-68) long; cheliceral blade 26 (23-30) long, curved with a single tooth distally.

Leg segmentation formula $fSp = 6-6-6$. IP = 1298 (1280-1320). Leg I 448 (431-456); trochanter 53 (50-55), with 1B; femur 83 (80-84), with 5B; genu 49 (47-50), with 4B, two solenidia σ , and a microseta κ ; tibia 78 (75-80), with 5B, two solenidia ϕ , and a microseta κ ; tarsus 104 (96-108), with 17B, one solenidion ω , one dorsal eupathidium ζ , one terminal eupathidium ζ , one famulus ϵ ; claw-like empodium 20 (19-21), two lateral claws 32 (32-34) each. Leg II 414 (406-419); trochanter 52 (49-55), with 1B; femur 74 (71-75), with 4B; genu 40 (38-42), with 3B, two solenidia σ and one microseta κ ; tibia 71 (70-72), with 5B and two solenidia ϕ ; tarsus 95 (92-96), with 14B, one solenidion ω , and one famulus ϵ ; claw-like empodium 22 (21-23), two lateral claws 31 (25-37) each. Leg III 436 (423-451); trochanter 57 (55-64), with 1B; femur 77 (75-79) with 4B; genu 42 (38-46), with 3B and two solenidia σ ; tibia 83 (80-86), with 5B; tarsus 102 (100-107), with 13B; claw-like empodium 26 (25-27), one lateral claw 37 (30-40).

Remarks

Larvae of 13 *Allothrombium* species have been recognized worldwide (Zhang and Xin 1992; Zhang & Norbakhsh 1995). Four species are known only from Europe: *A. fuliginosum* (Hermann), *A. recki* Feider & Agekian, *A. neapolitanum* Oudemans, and *A. mon-*


spessulanum Robaux & Aeschlimann (Feider 1951, Feider & Agekian 1967, Henking 1882, Hirst 1926, Robaux 1972, 1974, Robaux & Aeschlimann 1987, Oudemans 1910, 1912, Thor & Willmann 1947, Turk & Turk 1952). Six species have been recorded only from Asia: *A. ovatum* Zhang & Xin, *A. kekko* (Southcott), *A. epiphyllus* Shiba, and *A. chanaanense* Feider, *A. triticism* Zhang, and *A. mossi* Zhang (Feider 1977, Shiba 1976, Southcott 1986, Zhang & Xin 1992, Zhang & Norbakhsh 1995). Two species are known only from North America: *A. lerouxi* Moss and *A. mali* (Childers & Vercammen-Grandjean 1980; Moss 1962). *A. pulvinum* Ewing appears to be a cosmopolitan species and has been reported from Europe, Asia (China and Iran), and North America (Howard 1918, Miller 1925, Minks & Harrewijn 1988, Zhang 1988, 1991b, Zhang & Faraji 1994, Zhang & Xin 1989a, b, 1992), although its presence in Europe needs to be confirmed. The new species, *A. shirazicum*, is related to *A. triticism*, but can be distinguished from the latter from its long legs: leg lengths of I-II-III are 448-414-436 for *A. shirazicum*, but 335-328-367 for *A. triticism*.

Monotrombium simplicium Zhang

Description

Larvae with the following features: $fD = 2-2-6-4-4-2=20$; $fV = 2-2-2u-2 = 8$; $fcx = 2-1-1$; $fnTr = 1-1-1$; $fnFe = 5-4-4$; $fnGe = 4-3-3$; $fnTi = 5-5-5$; $fnTa = 17-14-13$; $fSol = I(0-2-2-1), II(0-1-2-1), III(0-1-0-0)$; $fk = I(1-1), II(1-0), III(0-0)$; $f\zeta = 2-0-0$; $fe = 1-1-0$; $fPp = 0-0-0-BNN2-BBNN\omega$; IP = 805; tarsus III with reduced inner claw. IP = 785 (754-814).

Larva. – Measurements are means of four specimens, with range in parentheses. Idiosoma 492-530 long, 280-300 wide. Standard measurements of scutum and scutellum as follows: AM 36 (35-38); AA 51 (48-54); AL 33 (25-39); AW 62 (59-65); MA 32 (31-34); PL 47 (45-47); PW 79 (74-83); AP 35 (34-35); S 48 (43-52); SB 50 (49-50); ASB 66 (63-68); PSB 43 (36-49); SD 109 (99-117); W 94 (81-104); HS 32 (25-36); LSS 68 (59-81); SL 42 (39-45); SS 25 (23-26). Ocular sclerite, 23 (22-25) long and 13 (11-13) wide; the anterior eye (diameter 9-11) larger than the posterior one (diameter 5-7). Palpal 52 (51-54) long. Adoral seta 6 long. Subcapitular setae 7 (6-7) long, 12 (11-12) apart at base. Cheliceral base 44 (42-45) long; cheliceral blade 12 (11-14) long. IP = 785 (754-814). Leg I 258 (250-262) long; coxa I 51 (45-57) long; trochanter 33 (32-34) long; femur 48 (47-49) long; genu 29 (27-31) long; tibia 42 (38-47) long; tarsus 57 (51-61) long; claw-like empodium 21 (19-22) long, two lateral claws 15 (13-16) long each. Leg II 253 (243-268) long; coxa II 58 (54-61); trochanter 32 (31-32) long; femur 45 (43-45) long; genu 24 (22-



Figs. 4-6. *Allothrombium shirazicum* Zhang sp. n. Holotype larva. – 4, first leg; 5, second leg; 6, third leg.

25) long; tibia 40 (36-43) long; tarsus 53 (49-58) long; claw-like empodium 23(21-23) long, two lateral claws 16 (14-17) each. Leg III 275 long; coxa III 55 (54-56) long; trochanter 35 (34-36) long; femur 49 (43-52) long; genu 28 (25-31) long; tibia 46 (42-50) long; tarsus 61 (53-68); claw-like empodium 25 (23-28), one lateral claw 17 (16-18).

Material. – Larvae (ZQZ 93-0128-2a, b, and c), IRAN, Shiraz, on wheat aphids, 14.iv.1992, N. Rastegari.

Remarks

This species was first described from larvae parasitic on aphids in Shahrkord, Iran (Zhang & Norbakhsh 1995). This is a new record of this species from Shiraz, Iran. The Shiraz specimens are almost identical to those from Shahrkord. A minor exception is that the subcapitular setae are longer and more narrowly spaced at base in the former than in the latter; 7 long, 12 apart at base in Shiraz specimens but 5.3 long, 13 apart at base in specimens from Shahrkord.

ACKNOWLEDGEMENTS

We are grateful to Mr. Graham duHeaume of the International Institute of Entomology for drawing the six figures in the paper. Space and facilities during this study were kindly provided by the Department of Entomology, The Natural History Museum, London.

REFERENCES

- Childers, C. C. & P. H. Vercammen-Grandjean, 1980. *Aphithrombium mali*, a new genus and species in the family Trombidiidae (Acari: Parasitengonae) parasitic on *Aphis pomi* De Geer. – Journal of the Kansas Entomological Society 53: 720-726.
- Eickwort, G. C., 1983. Potential use of mites as biological control agents of leaf-feeding insects. In M. A. Hoy, G. L. Cunningham & L. Knutson (eds), Biological control of pests by mites: 41-52. University of California Press/ANR Publishing Co., Oakland.
- Feider, Z., 1951. Un trombidiid dusman al puricilor de plante. – Academia Republicii Populare Romine Filialia Iasi Studii si Cercetari Stiintifice Biologie si Stiinte Agricole 2: 481-497.
- Feider, Z., 1977. Contribution à la connaissance des larves d'acariens du bassin oriental de la Méditerranée. – Israel Journal of Zoology 26: 100-113.
- Feider, Z. & H. Agekian, 1967. Un nouvel acarien parasite des pucerons. – Travaux de Museum d'Histoire Naturelle 'Grigore Antipa' 7: 71-80.
- Henking, H., 1882. Beiträge zur Anatomie, Entwicklungs-geschichte und Biologie von *Trombidium fuliginosum* Hermann. – Zeitschrift für Wissenschaftliche Zoologie 37: 533-663.
- Hirst, S., 1926. Note on the development of *Allothrombium fuliginosum* Hermann. – Journal of the Royal Microscopical Society 1926: 274-276.
- Howard, C. W., 1918. A preliminary report on the Trombidiidae of Minnesota. – Report State Entomologist of Minnesota 17: 111-144.
- Miller, E. A., 1925. An introductory study of the Acarina or mites of Ohio. – Ohio Agricultural Experiment Station Bulletin 386: 82-172.
- Minks, A. K. & P. Harrewijn, 1988. Aphids. Their biology, natural enemies and control. Vol. B. – Elsevier, Amsterdam, 364 pp.
- Moss, W. W., 1962. The immature stages of the red velvet mite *Allothrombium lerouxi* (Acari: Trombidiidae). – Annals of the Entomological Society of America 55: 295-303.
- Oudemans, A. C., 1910. Acarologische aantekeningen XXXI. – Entomologische Berichten, Amsterdam 3: 47-51.
- Oudemans, A. C., 1912. Die bis jetzt bekannte Larven von Trombidiidae und Erythraeidae mit besonderer Berücksichtigung der für den Menschen schädlichen Arten. – Zoologische Jahrbücher (Supplement) 14 (1) 230 pp.
- Robaux, P., 1972. Étude des larves de Trombidiidae; IV. Redescription des larvaires formes d'*Allothrombium neapolitum* Oudemans, 1910, *Neothrombium neglectum* (Bruyant), 1908 et *Microthrombium fasciatum* (Koch), 1836. – Acarologia 14: 612-630.
- Robaux, P., 1974. Recherches sur le développement et la biologie des acariens 'Trombidiidae'. – Memoires du Museum National d'Histoire Naturelle Serie A Zoologie 85: 1-186.
- Robaux, P. & J. P. Aeschlimann, 1987. *Allothrombium monspessulanum* nov. spec. (Acari: Trombidiidae), un important ennemi naturel des arthropodes inféodés à la luzerne cultivée (*Medicago sativa* Linnaeus) en région méditerranéenne. – Mitteilungen der Schweizerischen Entomologischen Gesellschaft 60: 43-50.
- Shiba, M., 1976. Taxonomic investigation on free-living Prostigmata from the Malay Peninsula. – Nature and life in Southeast Asia 7: 83-299.
- Southcott, R. V., 1986. Studies on the taxonomy and biology of the subfamily Trombidiidae (Acarina: Trombidiidae) with a critical revision of the genera. – Australian Journal of Zoology (Supplementary Series) No. 123: 1-116.
- Thor, S. & C. Willmann, 1947. Acarina. Trombidiidae. – Das Tierreich 71b: 187-541.
- Türk, F. A. & S. M. Türk, 1952. Studies on Acari - 7th Series. Records and descriptions of mites new to the British fauna together with short notes on the biology of sundry species. – Annals and Magazine of Natural History (12) 5: 497-500.
- Welbourn, W. C., 1983. Potential use of trombidioid and erythraeid mites as biological control agents of insect pests. In M. A. Hoy, G. L. Cunningham & L. Knutson (eds), Biological control of pests by mites. Agricultural Experiment Station. Division of Agriculture and Natural Resources. Special Publication 3304: 103-140. University of California.
- Welbourn, W. C. & O. P. Young, 1988. Mites parasitic on spiders, with a description of a new species of *Eutrombidium* (Acari: Eutrombidiidae). – Journal of Arachnology 16: 373-385.
- Zhang, Z.-Q., 1988. Two common mites of *Allothrombium* Berlese in China. – Kunchong Zhishi 25: 172-174 [in Chinese].

- Zhang, Z.-Q., 1991a. Biology of mites of Allothrombiinae (Acari: Trombidiidae) and their potential role in pest control. In F. Dusbabek & V. Bukva (eds), Modern acarology Vol.II.: 513-520. Academia, Prague.
- Zhang, Z.-Q., 1991b. Parasitism of *Acyrtosiphon pisum* (Harris) by *Allothrombium pulvinum* Ewing (Acariformes: Trombidiidae): Host attachment site, host size selection, superparasitism, and impact on host. – Experimental and Applied Acarology 11: 137-147.
- Zhang, Z.-Q. & F. Faraji, 1994. Notes on *Allothrombium pulvinum* Ewing (Acari: Trombidiidae) new to the fauna of Iran. – Acarologia 35: 357-360.
- Zhang, Z.-Q. & H. Norbakhsh, 1995. A new genus and three new species of mites (Acari: Trombidiidae) described from larvae ectoparasitic on aphids from Iran. – European Journal of Entomology 92: 705-718.
- Zhang, Z.-Q. & J.-L. Xin, 1989a. Biology of *Allothrombium pulvinum* Ewing (Acari: Trombidiidae), a potential biological agent of aphids in China. – Experimental and Applied Acarology 6: 101-108.
- Zhang, Z.-Q. & J.-L. Xin, 1989b. Studies on the morphology and life history of *Allothrombium pulvinum* Ewing (Acariformes: Trombidiidae). – Acta Entomologica Sinica 32: 192-199 [in Chinese with English abstract].
- Zhang, Z.-Q. & J.-L. Xin, 1992. A review of larval *Allothrombium* (Acari: Trombidiidae), with description of a new species ectoparasitic on aphids in China. – Journal of Natural History 26: 383-393.

Received 30 June 1995

Accepted 6 October 1995



Zhang, Zhi-Qiang and Rastegar-Pouyani, Nasrullah. 1996. "Larval mites (Acari: Trombidiidae) parasitic on aphids in Iran: key, a new species and new record." *Tijdschrift voor entomologie* 139, 91–96.

View This Item Online: <https://www.biodiversitylibrary.org/item/89566>

Permalink: <https://www.biodiversitylibrary.org/partpdf/65968>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.