

2. Descriptions of Two new Land-Shells from Madagascar and New Guinea. By HENRY ADAMS, F.L.S.

[Received April 27, 1875.]

(Plate XLV.)

EURYCRATERA FARAFANGA, sp. nov. (Plate XLV. figs. 1, 1a.)

E. testa imperforata, depresso-ovata, solida, rugis obliquis irregularibus sculpta, pallide fulva, fasciis plurimis rufo-castaneis, nonnullisque latis ornata; spira brevi, subconoidea; anfr. 4, convexiusculis, rapide accrescentibus, ultimo inflato; apertura obliqua, ampla, ovali, intus margaritacea, fasciis perlucentibus; perist. albo, incrassato, breviter expanso, marginibus callo crasso junctis, columellari dilatato.

Diam. maj. 60, min. 45, alt. 50 mill.

Hab. Madagascar; on a sandy plain in the south-west, near the Farafanga river.

PUPINOPSIS ANGASI, sp. nov. (Plate XLV. figs. 2, 2a.)

P. testa perforata, pupæformi, solida, rugoso-malleata, non nitente, rubida vel pallide fulva; spira medio tumida, in conum depressum terminata; anfr. 7, superis convexis, penultimo latere aperturæ planato, ultimo angustiore, antice descendente, basi juxta perforationem cristato; apertura verticali, circulari, bicanaliculata; perist. incrassato, flavido vel albido, reflexo, margine dextro sursum producto, canali levi a parietali separato, margine columellari medio inciso, canalem apertum extus dilatatum callo circumvallatum formante.

Long. 20, diam. 12 mill.

Hab. Louisiade archipelago, south-east of New Guinea.

This species is closely allied to *P. grandis*, Forbes, but differs from it in being very much smaller, in having a narrow umbilicus or perforation, and in the first four whorls of the spire being more depressed.

3. Descriptions of Three new Species of Shells from Australia.

By GEORGE FRENCH ANGAS, Corr. Mem. Z.S., F.L.S., &c.

[Received April 27, 1875.]

(Plate XLV.)

1. *HELIX FORRESTIANA*, n. sp. (Plate XLV. figs. 3, 3a.).

Shell narrowly perforate, globosely trochiform, rather thin, obliquely striated, very finely transversely granulated; from the apex to the middle of the last whorl fulvous chestnut, with an indication of a band of a lighter colour between the periphery and the suture of the last whorl, below the periphery pale yellowish brown; spire obtusely conical; whorls 5, slightly convex, the last rather inflated,

descending in front, a little contracted behind the aperture; aperture very oblique, ovate, purplish brown within; peristome not thickened, widely expanded, the margins approximating, joined by a thin callus; the right margin slightly flexuous, brown; the columellar margin white, dilated and reflexed, nearly concealing the perforation.

Diam. maj. 10, min. 7, alt. 7 lines.

Hab. North-west Australia.

This shell differs from the other Australian *Helices* to which it is allied, in having the outer lip thin and very widely expanded. I have named it after Mr. John Forrest, whose recent explorations in Western and Central Australia have added to our knowledge of a hitherto unknown portion of that island continent.

2. *HELIX BROUGHAMI*, n. sp. (Plate XLV. figs. 4, 4a.)

Shell narrowly perforate, conically globose, rather thin, obliquely striated and obscurely minutely granulated, very pale brown above, white below, with three narrow reddish brown bands—one at the suture, one above, and one below the periphery of the last whorl; spire obtusely and depressedly conical; whorls $5\frac{1}{2}$, slightly convex, the last rounded, descending in front; aperture oblique, circularly lunate; peristome slightly expanded and reflexed, the margins approximating, the columellar margin almost straight, dilated above, and nearly covering the perforation.

Diam. maj. 12, min. 10, alt. 10 lines.

Hab. Port Lincoln, South Australia.

Allied to *H. cassandra*, Pfr., but differs in having the spire considerably more elevated, and in the columellar margin being straightened and dilated and reflexed, nearly concealing the perforation.

3. *EURYTA BRAZIERI*, n. sp. (Plate XLV. figs. 5, 5a.)

Shell somewhat narrowly fusiform, the spire a little longer than the aperture, white, shining, ornamented with three broad transverse sharply defined bands of irregular descending olive-brown lines here and there passing into blotches, the central band the broadest; apex olive-brown; whorls 7, slightly convex, longitudinally plicate, the last whorl more narrowly and finely plicate, with the interstices crossed by fine liræ, the plicæ ceasing above the central band, faintly transversely sulcate at the base; aperture narrow; columella slightly subflexuous and covered with a thin callus; outer lip simple, acute.

Long. 6, lat. $1\frac{1}{2}$ lines.

Hab. Dredged off Port-Jackson Heads, in 25 fathoms.

A prettily marked shell, quite distinct from any of the other Australian species of this genus hitherto described.

DESCRIPTION OF PLATE XLV.

Figs. 1, 1a. *Eurycratera farafanga*, p. 389.

2, 2a. *Pupinopsis angasi*, p. 389.

3, 3a. *Helix forrestiana*, p. 389.

4, 4a. *Helix broughami*, p. 390.

5, 5a. *Euryta brazieri*, p. 390.



Angas, George French. 1875. "Descriptions of three new species of shells from Australia." *Proceedings of the Zoological Society of London* 1875, 389–390.

View This Item Online: <https://www.biodiversitylibrary.org/item/90418>

Permalink: <https://www.biodiversitylibrary.org/partpdf/67157>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.