

Sect. c. *Lamina of lateral wing broad, marginal ridge feebly developed; second to eleventh dorsal segments multisulcate.*

23. ZEPHRONIA SULCATULA.

Zephronia sulcatula, Butler, Ann. & Mag. Nat. Hist. ser. 4, vol. x. p. 357. n. 13, pl. 18. f. 5 (1872).

Borneo (*W. Jeakes*).

Type, B.M.

EXPLANATION OF PLATE XIX.

Fig. 1. *Sphærotherium glabrum*, profile.

1 a. — — —, anterior segments from above.

2. — — — *fraternum*, profile.

3. — — — *stigmaticum*, perspective view from behind.

4. *Zephronia noticeps*, head, nuchal plate, and anterior segments, enlarged; front view.

5. — — — *lutescens*, profile.

6. — — — *ignobilis*, profile, enlarged.

7. — — — *pilifera*, profile.

8. — — — *corrugata*, profile.

9. — — — *banksiana*, profile.

10. — — — *leopardina*, profile.

5. Descriptions of Eight new Species of Land and Marine Shells from various Localities. By GEORGE FRENCH ANGAS, Corr. Mem. Z.S., F.R.G.S., F.L.S., &c.

[Received December 16, 1872.]

(Plate XX.)

1. EUTHRIA ARACANENSIS, n. sp. (Plate XX. fig. 1.)

Shell ovately fusiform, solid, transversely ridged, the interstices finely transversely striate, the ridges stronger in front, pale olivaceous brown, ornamented with longitudinal irregular chestnut markings; whorls 9 to 10, strongly angulate and nodose in the middle, excavated above, and slightly rounded below; aperture ovate, nearly half the length of the shell, angulate posteriorly, pale flesh-colour; outer lip crenulate, finely sulcate within; columella strongly arcuate, with a moderate posterior callus; the rostrum slanting, recurved.

Length 2 inches, diam. 1 inch (*coll. Hanley*).

Hab. Aracan.

2. HELIX (SEMICORNU) SILENUS, n. sp. (Plate XX. fig. 2.)

Shell deeply and perspectively umbilicated, rather thin, globosely depressed, pale brown, minutely and very finely punctately striate; spire immersed; whorls 5, rounded, rapidly increasing, the last inflated, a little flattened at the upper part, descending considerably in front, and somewhat angulated round the umbilicus; aperture oblique, rhomboidly lunate; peristome thin, whitish; margins approximating, the right flexuous, narrowly expanded; the columellar margin rounded, expanded, and slightly reflexed.

Maj. diam. 7, min. 6; height 5 lines.

Hab. New Ireland.





Angas, George French. 1873. "Descriptions of eight new species of land and marine shells from various localities." *Proceedings of the Zoological Society of London* 1873, 182–184.

View This Item Online: <https://www.biodiversitylibrary.org/item/90422>

Permalink: <https://www.biodiversitylibrary.org/partpdf/67193>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.