depression; antennæ and lower part of face obscure ferruginous, the third joint of former not much longer than the second, the fourth joint the longest. Thorax with the anterior angles but slightly produced and rounded; surface finely punctured, shining testaceous or flavous. Elytra widened behind, rather deeply and very closely punctured, with a distinct longitudinal depression in the middle of the base; the latter narrowly ferruginous, which colour also extends in form of a longitudinal short streak down the shoulders to about one third the length of the elytra; another sutural semi-square spot is placed below the scutellum, while a transverse short fascia occupies the middle, and another similar-shaped spot is placed near the apex of each elytron. Neither of these markings touch the suture or the lateral margins. Claw-joint strongly inflated.

Hab. Irazu Mountain, Costa Rica, and Nicaragua.

16. ŒDIONYCHIS SEPTEMMACULATA, Sp. nov.

Piceous below; lower part of vertex, thorax, and elytra testaceous, the latter with seven black patches, viz. two at the base, one common to both elytra before the middle, the other four at the middle and before the apex respectively.

Length 3 lines.

Vertex impunctate, lower part of the latter, as well as that of the clypeus, testaceous; antennæ piceous, basal joints paler. Thorax rather convex, sides broadly margined, the anterior angles produced into a short tooth; surface minutely punctured, testaceous; scutellum obscure piceous: elytra narrowed at base and apex, broadly margined, with two short longitudinal depressions at their posterior half, more distinctly punctured than the thorax, of a light testaceous colour, with the patches of the following shape—the basal one obliquely cut at its posterior margin, the sutural one of triangular shape, followed closely by a transverse quadrate fascia, and the apical one also of a transversely subquadrate shape; none of these markings touch the sutural or the lateral margins. Claw-joint greatly dilated.

Hab. Peru.

2. On a Fourth Collection of Birds made by the Rev. G. Brown, C.M.Z.S., on Duke-of-York Island and in its vicinity. By P. L. Sclater, M.A., Ph.D., F.R.S., Secretary to the Society.

[Received May 2, 1879.]

(Plates XXXVI., XXXVII.)

I have now again the pleasure of laying before the Society a series of bird-skins collected by our excellent correspondent Mr. Brown on Duke-of-York Island and on the adjacent parts of New Britain and New Ireland. Though many of these specimens belong to species







J.Smit lith.

Hanhart imp.

1.MYZOMELA CINERACEA. 2.DONACICOLA SPECTABILIS.

3. MUNIA FORBESI.



represented in the former collections1, there are several novel and interesting birds in the present consignment, to which I wish to draw special attention, and which show that the avifauna of this district

is by no means yet completely known to us.

The total number of skins in the present Collection, including some just previously received in a letter, is 59, belonging to 41 species. Every skin being labelled with an exact date and locality, it will be advisable to give a complete list of the species.

* 1.	Graucalus sclateri, Salvad	New Ireland.
* 2.	Grancalus sublineatus, sp. nov	Topaia, New Ireland.
3.	Dicrurus læmostictus, Scl	New Britain.
* 4.	Pachycephala melanura, Gould	Palakuüru Island.
* 5.	Myzomela sclateri, Forbes	Palakuüru Island.
* 6.	Myzomela cineracea, sp. nov	New Britain.
*7.	Myzomela erythrina, Ramsay	New Ireland.
8.	Philemon cockerelli, Scl	New Britain.
* 9.	Donacicola spectabilis, sp. nov	New Britain.
*10.	Munia forbesi, sp. nov	Topaia, New Ireland.
11.	Dendrochelidon mystacea (Less.)	Duke-of-York group.
*12.	Collocalia, sp. inc	Duke-of-York group.
*13.	Collocalia, sp. inc	New Britain.
*14.	Eurystopodus sp. inc. (pullus)	Topaia, New Ireland.
*15.	Alcyone pusilla (Temm.)	Wood Harbour, New Ireland.
16.	Halcyon sanctus, Vig. and Horsf	Duke-of-York group.
*17.	Scythrops novæ-hollandiæ	Duke-of-York group.
18.	Scythrops novæ-hollandiæ Cacomantis sp. inc. (jr.)	Duke-of-York group.
*19.	Chrysococcyx lucidus (Gm.)	New Ireland.
20.	Lorius hypænochrous, Gray	New Britain.
91	Trichoalossus massenæ Bn	New Ireland.
22.	Ninox variegata (Q. et G.)	Topaia, New Ireland.
*23.	Ninox jacquinoti, Bp	New Dritain.
*24.	Henicopernis longicauda (Garn.)	New Britain.
25.	Baza reinwardti (Müll. et Scl.)	New Ireland.
*26.	Falco lunulatus, Lath	Duke-of-York group.
*27.	Astur torquatus (Temm.)	New Ireland.
28.	Astur etorques (Salvad.)	New Britain.
29.	Carpophaga van-wycki, Cassin	Duke-of-York group.
30.	Œdirhinus insolitus (Schl.)	Duke-oi-York group.
31.	Phlogenas johannæ, Sclater	Duke-of-York group.
32.	Macropygia carteretia, Bp	New Britain and Duke-of-York
		group.
33.	Macropygia nigrirostris, Salvad	Dake-of-York group.
*34.	Excalfactoria chinensis (Linn.)	Duke-of-York group.
*35.	Amaurornis moluccana (Wallace)	Duke-oi-1 ork group.
36	Charadrius fulvus (Gm.)	Makado, Duke-of-York group.
37.	Ægialitis dubia (Scop.)	Topaia, New Ireland.
38.	Numenius uropygialis, Gould	Duke-or-rork group.
39.	Totanus ineanus (Gm.)	Duke-of-York group.
40.	Tringoides hypoleucus (Linn.)	Duke-of-York group.
41.	Sterna fuliginosa (Gm.)	Duke-of-York group.
	The species thus marked have not been re	
	The species that marked have hot been a	1

collections.

1. Graucalus sclateri, Salvad.; Sharpe, Cat. iv. p. 40.

A female of this species, differing from the type (now in the Bremen Museum), which is doubtless a male, in its ashy grey lores See P. Z. S. 1877, p. 96, 1878, p. 289, and 1878, p. 670.



1879. "On a Fourth Collection of Birds made by the Rev. G. Brown on Duke of York Island and its vicinity." *Proceedings of the Zoological Society of London* 1879, 446–451. https://doi.org/10.1111/j.1096-3642.1879.tb02677.x.

View This Item Online: https://www.biodiversitylibrary.org/item/90455

DOI: https://doi.org/10.1111/j.1096-3642.1879.tb02677.x

Permalink: https://www.biodiversitylibrary.org/partpdf/67260

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.