

A NEW GENUS OF PRIMITIVE CRANE FLIES (DIPTERA: TANYDERIDAE) IN CRETACEOUS BURMESE AMBER, WITH A SUMMARY OF FOSSIL TANYDERIDS

GEORGE POINAR, JR. AND ALEX E. BROWN

(GP) Department of Zoology, Oregon State University, Corvallis, OR 97331, U.S.A.
(e-mail: poinarg@casco.net); (AEB) 629 Euclid Avenue, Berkeley, CA 94708, U.S.A.

Abstract.—A new genus and species of primitive crane flies, *Dacochile microsoma* Poinar and Brown (Tanyderidae) is described from Cretaceous Burmese amber. It differs from extant and extinct members of the family by the following combination of characters: small size (wing length, 2.8 mm), reduced anal lobe, hyaline wing membrane, crossvein $cua-a_1$ forming cell cua , very short vein R_{2+3} , very long terminal maxillary palps, and mandibles. The well-developed mandibles indicate that the species obtained food by piercing and sucking. A list of fossil tanyderids is presented.

Key Words: *Dacochile* n. gen., *Dacochile microsoma* n. sp., Tanyderidae, Burmese amber, Cretaceous


The Tanyderidae, or primitive crane flies, possess many plesiomorphic characters, especially regarding the wing venation. The family consists of 12 extant genera, most in the Southern Hemisphere. The adults occur in moist terrestrial habitats while the larvae are found in shallow water or along the margins of lakes, rivers and streams, usually in sediment or rotting vegetation (Alexander 1981). Origin of the family and its relationship with other Nematocera has been of interest, and fossils from the Eocene and Jurassic have been reported (Table 1).

A female tanyderid discovered in Cretaceous Burmese amber, described below in a new genus, provides insight into the morphology of early members of this interesting group. Amber from Burma occurs in lignitic seams in sandstone-limestone deposits in the Hukawng Valley (Poinar 1992, Zherikhin and Ross 2000). Nuclear magnetic resonance (NMR) spectra of amber samples taken from the same locality as the

fossil indicated an araucarian (possibly *Agathis*) source of the amber (Lambert and Wu, unpublished research 2002). Palynomorphs obtained from the amber beds where the fossil originated have been assigned to the Upper Albian (~100–110 mya) of the Lower Cretaceous (Cruickshank and Ko 2002).

MATERIALS AND METHODS

The amber was recut and polished in order to view the specimen better. The amber piece containing the fossil is rectangular in outline, measuring 7 mm in greatest length, 5 mm in greatest width and 1 mm in greatest depth. The specimen is well preserved and nearly complete, with only the tips of the tibiae and ovipositor missing. Observations, drawings and photographs were made with a Nikon SMZ-10 stereoscopic microscope and Nikon Optiphot optical microscope (with magnifications up to 650 $\times$). In the following description, terminology is



Figs. 1–3. *Dacochile microsoma*. 1, Holotype in Burmese amber. Bar = 836 μm . 2, Proboscis with well-developed mandibles (arrows). Bar = 95 μm . 3, Anal region of wing showing long setae emerging from costal vein (arrow). Note microtrichia on wing membrane. Bar = 103 μm .

that presented in the "Manual of Nearctic Diptera" (McAlpine 1981).

The presence of a radius with 5 branches all ending at the wing margin, the subcosta bifurcate at its extremity, with the basal branch appearing as a crossvein ending in R_1 , a single anal vein reaching the margin, the costa continuing around the wing margin, setae between the ommatidia and cell br (first basal cell) longer than cell bm (second basal cell) place the fossil in the family Tanyderidae Osten-Sacken 1879.

***Dacochile* Poinar and Brown,
new genus**

Type species: Dacochile microsoma Poinar and Brown.

Description.—Body (length, 2.6 mm) and wing (length, 2.8 mm) small, anal lobe greatly reduced, anal margin slightly concave; wing membrane hyaline, crossvein cu a-a₁ forming cell cu a, vein R_{2+3} short, mandibles well developed.

Etymology.—Daco is from the Greek "dakno" for bite; chile is from the Greek "chilo" for lip; the gender is neuter.


Diagnosis.—In Williams' (1933) key to the extant and fossil Tanyderidae, *D. microsoma* would align with the Baltic amber fossil *Macrochile* Loew in the couplet "wings immaculate." Whether the wings of *Macrochile* are completely hyaline is questionable since Alexander (1931) noted vague indications of dusky spots at the end of vein Sc, on r-m and at the fork of M3 and CuA1 on *M. spectrum* Loew, and Podenas (1997) noted dark areas on the wings of *M. baltica* Podenas. The two described species of *Macrochile* are large (body length from 8.2 mm to 12 mm and wing length from 4.5 mm to 8.5 mm) and have distinct anal lobes. The Rs varies from shorter to approximately the same length as R_{2+3} in *Macrochile* while it is 3 times as long as R_{2+3} in *Dacochile*. The extant and extinct Jurassic species of *Protanyderus* Handlirsch differ from *Dacochile* in having pictured wings with an anal lobe and in lacking the cu a-a₁ crossvein. The presence

of a distinct cu a-a₁ crossvein, forming a cu a cell is rare in tanyderids. However, it occurs in the Patagonian species *Neoderus patagonicus* Alexander (Williams 1933).

The extinct genus *Praemacrochile* Kalugina differs from *Dacochile* in having Sc end opposite the middle of R_{4+5} , R_2 short, almost $\frac{1}{3}$ of Sc and $\frac{1}{3}$ of R_{2+3} and having R_{4+5} as long as or slightly longer than R_2 . *Dacochile* shares some characters with *Nannotanyderus* Ansorge 1994, namely the small size and reduced anal lobe, but the latter genus is characterized by a short Rs vein, only half as long as R_{2+3} , a large cell bm and a rounded anal lobe (Ansorge 1994, Ansorge and Krzemiński 2002).

***Dacochile microsoma* Poinar and
Brown, new species**
(Figs. 1–8)


Description.—Holotype female; with characters listed under the generic diagnosis. General coloration brown; body length 2.6 mm. **Head:** Length: 315 μ m (not including rostrum), eyes dichoptic, with short hairs arising between facets of ommatidia; antenna 17-segmented, 2.37 mm long, segment 1: 63 μ m, 2: 113 μ m, 3: 63 μ m, 4: 283 μ m, 5: 214 μ m, 6: 189 μ m, 7: 189 μ m, 8: 145 μ m, 9: 170 μ m, 10: 95 μ m, 11: 107 μ m, 12: 107 μ m, 13: 101 μ m, 14: 107 μ m, 15: 107 μ m, 16: 107 μ m, 17: 107 μ m. Antennal segments with numerous uniform setae; conspicuous verticils equal or exceed corresponding segments in length, apical segment as long as preceding segment, postpedicel short, equal in length to pedicel; maxillary palp 5-segmented, 850 μ m long, with 5th segment 2.6 times as long as 4th, segment 1: 50 μ m, 2: 132 μ m, 3: 145 μ m, 4: 145 μ m, 5: 378 μ m; head bearing distinct, short rostrum, 265 μ m long, with strong, pointed mandibles lacking noticeable teeth on inner surface (details shown in Figs. 2, 7). **Thorax:** Brown, 790 μ m long; legs yellowish brown, coxa 1: 334 μ m, 2: 410 μ m, 3: 422 μ m, trochanter 1: 113 μ m, 2: 126 μ m, 3: 113 μ m, femur 1: 1.28 mm, 2: 1.43 mm, 3: 1.42 mm; all tarsi and tips



Figs. 4-5. *Dacochile microsoma*. 4, Flagellomere showing verticils at the base of each segment and dense setae uniformly covering each segment. Bar = 89 μm . 5, Extremely long terminal segment of maxillary palp. Bar = 48 μm .

of tibiae missing; wing hyaline, all veins bearing setae, long setae on costa in anal area, wing membrane covered with microtrichia, Sc relatively short, less than half wing length, ending before origin of R_{2+3} ; cell br over twice as long as cell bm; crossvein m-cu absent; Cu P vein faint, arising

from A_1 and curving toward wing margin; base of Cu A_1 appears as crossvein; tip of veins R_1 , R_2 and R_3 bent anteriorly at wing margin; anal lobe greatly reduced or possibly absent, replaced with long setae; halter brownish yellow, 258 μm long, covered with short setae; further details of venation



Figs. 6–8. *Dacochile microsoma*. 6, Wing showing veins and cells referred to in this study. Bar = 627 μ m. Setae and microtrichia omitted. 7, Lateral view of head. l = labrum, la = labium, lb = labellum, m = mandible. Bar = 78 μ m. 8, Lateral view of thorax. ke = katepimeron + meron, kt = katepisternum, m = metanotum, mk = metakatepisternum, p = prosternum, ps = postscutellum, s = scutellum, sc = scutum. Bar = 165 μ m.

Table 1. Fossil species of Tanyderidae.

Species	Locality	Age	Reference
<i>Dacochile microsoma</i> , n. sp.	Burma	Cretaceous	present work
<i>Macrochile baltica</i> Podenas	Baltic amber	Eocene	Podenas 1997
<i>Macrochile spectrum</i> Loew	Baltic amber	Eocene	Loew 1850
<i>Nannotanyderus grimmenensis</i> A. & K.	Germany	Lower Jurassic	Ansorge & Krzemiński 2002
<i>Nannotanyderus krzeminskii</i> Ansorge	Germany	Lower Jurassic	Ansorge 1994
<i>Praemacrochile chinensis</i> K. & R.	China	Middle Jurassic	Krzemiński & Ren 2001
<i>Praemacrochile decipiens</i> (Bode)	Germany	Lower Jurassic	Ansorge & Krzemiński 2002
<i>Praemacrochile dobbertinensis</i> A. & K.	Germany	Lower Jurassic	Ansorge & Krzemiński 2002
<i>Praemacrochile stackelbergi</i> Kalugina	Siberia	Upper Jurassic	Kalugina & Kovalev 1985
<i>Protanyderus mesozoicus</i> Kalugina	Mongolia	Upper Jurassic	Kalugina 1988
<i>Protanyderus senilis</i> Kalugina	Mongolia	Upper Jurassic	Kalugina 1992
<i>Protanyderus vetus</i> Kalugina	Mongolia	Upper Jurassic	Kalugina 1992

shown in Fig. 6 and of thorax in Fig. 8. *Abdomen*: Brown, extended, 1.64 mm long, twisted unnaturally about 90° at approximately ⅔ of its length, tip of ovipositor missing.

Male.—Unknown.

Material examined.—Holotype female in Burmese amber, deposited in the Poinar amber collection (accession # B-D-12) maintained at Oregon State University.

Etymology.—Micro is from the Greek “mikros” for small and somus is from the Greek “soma” for body.

Diagnosis.—*Dacochile microsoma* is one of the smallest known members of the family. Other distinguishing characters are the short antennal postpedicel, the very short R_{2+3} and the extremely short bm cell. While members of the genus *Radinoderus* also have a relatively short R_{2+3} , they lack a cu-a cell, possess an anal lobe and have pictured wings. Most tanyderids have the terminal flagellomere shorter than the penultimate, and the terminal maxillary palp not more than twice the penultimate one (Williams 1933, Alexander 1981). However the terminal and penultimate flagellomeres are equal in *D. microsoma* and the terminal maxillary palp is 2.6 times longer than the penultimate. Unfortunately, head structures are not noted for the other Mesozoic fossil tanyderids (Table 1).

DISCUSSION

The presence of well-developed mandibles in *D. microsoma* is interesting because these structures have only been reported in the Australian genus *Radinoderus* Handlirsch and the North American *Protoplasa* Osten-Sacken (Downs and Colless 1967). In *Dacochile*, the size of the mandibles suggests that a piercing-sucking habit provided the major means of obtaining nourishment. The blood-sucking habit is well established in the Diptera, with each group evolving their own specialized type of mouthpart modification in order to obtain the food source (Downs 1970, Kalugina 1991).

It is interesting that the mouth structures closely resemble those of the Psychodidae (Quate and Vockeroth 1981). The similarity between the wing structure of Mesozoic Tanyderidae and Psychodidae has been noted, and the similarity of the mouth structure reported here supports the contention that the two families are closely related and may have originated from a common stem (Crampton 1926, Krzemiński and Evenhuis 2000).

A list of described fossil tanyderids is presented in Table 1. Twelve fossil species in 5 genera (one extant and 4 extinct) have been described, all from Old World deposits. The oldest members are from the Lower Jurassic and the youngest from the Eocene.

Only three species have been described from amber deposits. It is interesting that some of the Upper Jurassic forms have been placed in the extant genus *Protanyderus* Handlirsch. Since the Jurassic fossils were identified only on the basis of wing characters, we feel that their placement in an extant genus is premature and should be made only after an examination of body characters. If confirmed, these would represent the oldest terrestrial generic lineages, thus far limited to four insect genera found in Cretaceous Lebanese amber (Poinar and Milki 2001).

ACKNOWLEDGMENTS

We thank Neal Evenhuis, Jörg Ansorge, and Wiesław Krzemiński for supplying pertinent literature on fossil tanyderids and Roberta Poinar for comments on an early draft of the manuscript.

LITERATURE CITED

- Alexander, C. P. 1931. Crane-flies of the Baltic amber (Diptera). *Bernstein-Forschung* 2: 1–135.
- . 1981. Tanyderidae, pp. 149–151. In McAlpine, J. F. et al., eds. *Manual of Nearctic Diptera*. Monograph 27, Vol. 1. Agriculture Canada, Research Branch, Ottawa.
- Ansorge, J. 1994. Tanyderidae and Psychodidae (Insecta: Diptera) from the Lower Jurassic of north-eastern Germany. *Paläontologische Zeitschrift* 68: 199–210.
- Ansorge, J. and W. Krzemiński. 2002. Lower Jurassic tanyderids (Diptera: Tanyderidae) from Germany. *Studia Dipterologica* 9: 21–29.
- Crampton, G. C. 1926. The external anatomy of the primitive tanyderid dipteran *Macrochile spectrum* Loew, preserved in Baltic amber. *Bulletin of the Brooklyn Entomological Society* 21: 1–14.
- Cruikshank, R. D. and K. Ko. 2002. Geology of an amber locality in the Hukawng Valley, northern Myanmar. *Journal of Asian Earth Sciences* 21: 441–445.
- Downs, J. A. 1970. The ecology of blood-sucking Diptera: An evolutionary perspective, pp. 232–258. In Fallis, A. M., ed. *Ecology and Physiology of Parasites*, University of Toronto Press, Toronto.
- Downs, J. A. and D. H. Colless. 1967. Mouthparts of the biting and blood-sucking type in Tanyderidae and Chironomidae (Diptera). *Nature* 214: 1355–1366.
- Kalugina, N. S. 1988. Mesozoic Diptera Psychodomorpha and Tipulomorpha (Tanyderidae, Eoptychopteridae, Limoniidae: Diptera), pp. 81–88. In Rozanov, A. Yu., ed. *New Species of Fossil Invertebrates of Mongolia*. *Trudy Sovetsko-Mongolskaya paleontologeskaia Ekspeditsiya* 33. (In Russian.)
- . 1991. New Mesozoic Simuliidae and Leptoconopidae and the origin of bloodsucking in the lower dipteran insects. *Paleontological Journal* 25: 66–77.
- . 1992. Psychodomorph Dipterans from the Jurassic of the Mongolian Altai (Diptera: Tanyderidae, Eoptychopteridae). *Paleontological Journal* 26: 142–146.
- Kalugina, N. S. and V. G. Kovalev. 1985. Dipterous insects of Jurassic Siberia. *Paleontological Institute, Akademia Nauk, Moscow*, 198 pp. (In Russian.)
- Krzemiński, W. and N. L. Evenhuis. 2000. Review of Diptera palaeontological records, pp. 535–564. In Papp, L. and B. Darvas, eds. *Manual of Palearctic Diptera*. Science Herald, Budapest, Vol. 1.
- Krzemiński, W. and D. Ren. 2001. *Praemacrochile chinensis* sp. n. from the Middle Jurassic of China (Diptera: Tanyderidae). *Polskie Pismo Entomologica* 70: 127–129.
- Loew, H. 1850. *Ueber den Bernstein und die Bernsteinfauna*. E. S. Mittler & Sohn, Berlin, 44 pp.
- McAlpine, J. F. 1981. Morphology and terminology—Adults, pp. 9–63. In McAlpine, J. F. et al., eds. *Manual of Nearctic Diptera*. Monograph 27, Vol. 1, 674 pp. Agriculture Canada, Research Branch, Ottawa.
- Podenas, S. 1997. New *Macrochile* Loew, 1850 (Diptera, Tanyderidae) from the Baltic amber. *Mitteilungen der Geologisch-Paläontologischen Institut der Universität Hamburg* 80: 173–177.
- Poinar, Jr., G. O. 1992. *Life in Amber*. Stanford University Press, Palo Alto, 350 pp.
- Poinar, Jr., G. O. and R. Milki. 2001. Lebanese amber. The oldest insect ecosystem in fossilized resin. Oregon State University Press, Corvallis, 97 pp.
- Quate, L. W. and J. R. Vockeroth. Psychodidae, pp. 153–190. In McAlpine, J. F. et al., eds. *Manual of Nearctic Diptera*. Monograph 27, Vol. 1, Agriculture Canada, Research Branch, Ottawa.
- Williams, I. 1933. The external morphology of the primitive tanyderid dipteran *Protoplasma fitchii* O.S., with notes on the other Tanyderidae. *Journal of the New York Entomological Society* 41: 1–36.
- Zherikhin, V. V. and A. J. Ross. 2000. A review of the history, geology and age of Burmese amber (Burmite). *Bulletin of the Natural History Museum (London) (Geology)* 56: 3–10.



Poinar, George O. and Brown, Alex E. 2004. "A new genus of primitive crane flies (Diptera: Tanyderidae) in Cretaceous Burmese amber, with a summary of fossil tanyderids." *Proceedings of the Entomological Society of Washington* 106, 339–345.

View This Item Online: <https://www.biodiversitylibrary.org/item/95279>

Permalink: <https://www.biodiversitylibrary.org/partpdf/68805>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Entomological Society of Washington

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.