February 15, 1887.

Prof. W. H. Flower, LL.D., F.R.S., President, in the Chair.

The Secretary read the following report on the additions to the Society's Menagerie during the month of January 1887:—

The total number of registered additions to the Society's Menagerie during the month of January was 38. Of these 6 were by birth, 21 by presentation, 5 by purchase, 1 by exchange, and 5 were received on deposit. The total number of departures during the same period, by death and removals, was 119.

The most noticeable additions during the month were:—

1. Two Blakiston's Owls (*Bubo blakistoni*) from Japan, presented by J. H. Leech, Esq., F.Z.S., January 20th, new to the Society's Collection.

Mr. Leech informs me that he procured the two specimens in question from Mr. H. Henson of Hakodate, Yesso. Mr. Henson had bought them from a native hunter, who took them for young eagles, which are common in Yesso. Their exact locality is the lakes 20 miles north of Hakodate, and this is a new locality for the species, of which, Mr. Leech believes, only four specimens were previously known¹.

Mr. J. H. Gurney, F.Z.S., has kindly furnished me with the following notes upon these birds:—

"Blakiston's Eagle-Owl appears to belong to the Bubonine genus (or subgenus) *Pseudoptynx* of Kaup, instituted by that author for the reception of *P. philippensis* (Gray), and readily distinguished from *Bubo* in having the toes bare, although the tarsi are feathered.

"*P. philippensis* is a native of the island of Luzon, from which I have seen specimens, and was figured by the late Lord Tweeddale in the Society's 'Transactions,' vol. ix. pl. xxv. fig. 2.

"The only other species of the genus is *P. gurneyi*, Tweed., from the island of Mindanao, which was figured by Lord Tweeddale in the 'Proceedings' of the Zoological Society, 1878, pl. lviii.

"The localities inhabited by the three species of the genus *Pseudoptynx* appear to indicate that they form a natural group, geographically as well as structurally."—J. H. G., Feb. 26, 1887.

2. Three Hooker's Sea-lions (*Otaria hookeri*), presented by the Hon. W. J. M. Larnach, C.M.G., Minister of Marine of New Zealand, received the 26th January.

Sir F. D. Bell, the Agent-General for New Zealand, informs me that these animals, which were captured at the Auckland Islands by Capt. John Fairchild, Master of the New-Zealand Government steamer 'Hinemoa,' were originally four in number (two males and two females), but that one died on the voyage.

It is very difficult to settle the species of *Otaria* without reference to the form of their palates and dentition; but, judging from the

¹ Cf. Seebohm, 'Ibis,' 1884, pp. 42, 183, pl. vi.; id. P. Z. S. 1883, p. 466.

Flower, William Henry. 1887. "February 15, 1887." *Proceedings of the Zoological Society of London* 1887, 138–139.

<https://doi.org/10.1111/j.1096-3642.1887.tb02952.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/99199>

DOI: <https://doi.org/10.1111/j.1096-3642.1887.tb02952.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/72990>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.