

CLYTANTHUS IGNOBILIS.

Cl. 4-punctato (Fabr.) *proxime affinis*. *Nigro-fuscus, tomento cinereo-flavo vestitus; elytris utrinque maculis 5 nigris, quam in Cl. 4-punctato majoribus, scilicet 1 curvata post scutellum, 1 parva humerali, 1 antico-discoïdali, 1 mediana majore transversa et 1 huic proxima longitudinali oblonga.*

Long. 6 lin.

Murree.

11. On a new Species of *Notauges* (*N. hildebrandti*, Cab.).

By Dr. G. HARTLAUB.

[Received June 17, 1878.]

The Museum of Natural History of Bremen has been enriched of late with a fine adult specimen of a new typical *Notauges*, discovered by the famous traveller Mr. Hildebrandt, in Eastern Africa. The exact place where he collected this interesting bird is "Tkanga in Ukamba."

Mas ad. *capite, gutture, dorso, tergo, uropygio, supracaudalibus, tectricibus alarum minoribus margineque alari chalybeis, nitore purpurascente; collo postico late et circumscripte aeneo-virescente; alis duabus seriebus macularum holosericeo-nigrarum bifasciatis; tectricibus alarum majoribus viridi-aeneis; remigibus primariis in pogonio externo aeneo-cærulescentibus, obscure maculatis, interno nigris; tectricibus minoribus in pogonio externo unicoloribus aeneo-virentibus; pectore, abdomine, cruribus et subcaudalibus cinnamomeo-rufis; subalaribus majoribus rufis, minoribus chalybeis; rectricibus cærulescenti-aeneis, minus distincte undulatis; rostro et pedibus nigris; iride aurantiaco-rubra.*

Long. 19 centim., rostr. a fr. 18 mill., al. 123 mill., caud. 83 mill., tars. 30 mill.

The steel-blue colour of the head is less brilliant on the crown; there appear some rufous feathers on the uropygium; and some of the upper tail-coverts have a slight rufous terminal edging; the first primary is of a uniform blackish colour.

In *Notauges superbus* (Abyssinia, Shoa) the colouring is very different: the upper parts are brown-green, the vent and under tail-coverts are white, and there are other differences.

Mr. Hildebrandt told me that during his many excursions in East Africa he never met with *Notauges albicapillus*, Blyth. This is certainly one of the rarest birds, the Calcutta specimen collected by Speke and Burton being as yet the only one known.



Harthlaub, G. 1878. "On anew Species of Notauges (N. hildebrandti, Cab.)."
Proceedings of the Zoological Society of London 1878, 721–721.
<https://doi.org/10.1111/j.1469-7998.1878.tb08011.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/90451>

DOI: <https://doi.org/10.1111/j.1469-7998.1878.tb08011.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/73526>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.