6. On a new Species of *Tringa* from Alaska.

By J. E. HARTING, F.L.S., F.Z.S.

[Received March 26, 1874.]

(Plate XL.)

Through the kindness of Professor Spencer Baird, I received some months ago a specimen of a Sandpiper from St. Paul's Island, Alaska, with a request that I would examine and report upon it. It resembled at first sight a very large Dunlin, *Tringa alpina*, in partial summer plumage, and with the breast more or less spotted with black; but its superior size showed at once that it could not belong to that species. The only other *Tringa* at all resembling it with which I was then acquainted being *Tringa crassirostris* of Temminck and Schlegel, from China, Japan, the Malay countries, and Australia, I hastily but erroneously came to the conclusion that it should be referred to that species; and without waiting to institute any comparison of specimens, I wrote to Professor Baird accordingly. This was unfortunate; for on subsequently making a more careful examination, and comparing the bird in question with specimens in my collection of both *T. alpina* and *T. crassirostris*, I found to my surprise that it differed materially from both, being much smaller than *T. crassirostris* although considerably larger than *T. alpina*, and in several other respects, as I shall presently point out, holding an intermediate position between these two species. I have now no hesitation in saying that it may be regarded as a new and hitherto undescribed bird, and I accordingly propose to name it *Tringa gracilis*.

It may be described as follows:—

TRINGA GRACILIS, sp. nov.

T. similis alpinæ sed conspicue major. Notæi plumis nigris, late rufo-marginatis; pileo fuscescente, rufonigroque striolato; capitis et colli lateribus dilute rufescentibus, maculis minutis fuscis; uropygio nigro; gula et fronte albis; macula pectorali magna nigra; abdomine crissoque albis; tectricibus alarum pallide fuscis, albido limbatis; remigibus fuscis, scapis pure albis; subalaribus albis; rectricibus lateralibus pallide fuscis, albo-limbatis; rostro et pedibus nigricantibus. (Ptil. æstiv.) Long. tot. 10 poll., rostr. 1·5, alæ 5·5, tars. 1, dig. med. cum ung. 1·1.

Hab. St. Paul's Island, Alaska.

The specimen from which the above description is taken was most kindly presented to me by Professor Baird, with the information that it had been obtained with several others on the island above mentioned in the month of July 1872. It is evidently in summer plumage; and being at this season black-breasted, like the Dunlin, *Tringa alpina*, we may fairly assume that in winter, like that species,

J G Keulemans lith.

1.2.3. TRINGA ALPINA. 4.5.6.T. GRACILIS. 7.8.9.T. CRASSIROSTRIS.

M & N Hanhart imp.

Harting, James Edmund. 1874. "On a new Species of Tringa from Alaska."
Proceedings of the Zoological Society of London 1874, 242–244.
<https://doi.org/10.1111/j.1096-3642.1874.tb02478.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/90419>

DOI: <https://doi.org/10.1111/j.1096-3642.1874.tb02478.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/73669>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.