

STUDIES IN THE LIABEAE (ASTERACEAE). XII.

A NEW SPECIES OF MUNNOZIA FROM

COSTA RICA

Harold Robinson

Department of Botany

Smithsonian Institution, Washington, DC. 20560.

The genus Munnozia contains about forty species which are mostly restricted to South America. The genus has previously been known in Central America only on the basis of the widely distributed M. senecionidis Benth. which extends its range into Costa Rica and western Panama. Recent collections have unexpectedly revealed a thoroughly distinct undescribed member of the genus that seems endemic to central Costa Rica. The specimens are the result of the collecting program of Dr. R. L. Wilbur of Duke University and the species is here named in his honor.



Munnozia wilburii H. Robinson, sp. nov.

Plantae suffrutescentes usque ad 2 m altae. Caules hexagonales hirsuti et appresse aliquantum evanescentiter albo-arachnoideo-tomentosi, nodis in alis petiolarum connatis inclusis. Folia opposita, petiolis ad 8 cm longis distincte alatis, alis inferne ad 16 mm latis superne angustioribus; laminae deltoideo-ovatae plerumque 9-16 cm longae et 5-12 cm latae base truncatae margine multo dentatae et mucronate denticulatae apice breviter acuminatae supra hirtellae subtus dense albe persistentiter arachnoideo-tomentosae, nervis secundariis basilaribus congestis in partibus trinervatis vel subtrinervatis. Inflorescentiae terminales profuse laxae pyramidaliter paniculatae inferne opposite et superne plerumque alternate ramosae, pedunculis 4-10 cm longis sparse pallide hirtellis et dense albo-arachnoideo-tomentosis. Capitula plerumque ca. 10 mm alta et 10-12 mm lata. Squamae involucris 30-35 late ovatae vel late oblongae distincte inaequales 4-7 mm longae et plerumque 2-3 mm latae apice obtusae vel breviter acutae extus glabrescentes 5-10 striatae inferne subcarnosae superne breviter herbaceae, bracteis exteriores minimis persistentiter sparse arachnoideo-tomentosae; receptacula fimbriis squamiformibus laciniatis ad 2 mm longis. Flores radii 40-45; corollae flavae, tubis 4-6 mm longis plerumque dense puberulis,

limbis 12-17 mm longis ad 2 mm latis base dense pilosis superne glandulosis et sparse puberulis. Flores disci 40-50; corollae flavae 7-8 mm longae, tubis 4-5 mm longis dense puberulis, faucis ca. 1 mm longis abrupte late campanulatis base sparse puberulis, lobis ca. 2 mm longis 0.8-1.0 mm latis extus multi-glanduliferis distaliter in fasciculo dense pilosis; filamenta in parte superiore ca. 0.3 mm longa; thecae nigrescentes ca. 2 mm longae, cellulis endothecialibus solum in parietibus transversalibus noduliferis; appendices antherarum oblongae ca. 0.3 mm longae et 0.2 mm latae. Achaenia ca. 1.3 mm longa longe setifera ca. 10-costata; setae pappi longiores 35-40 plerumque ca. 5 mm longae subtiliter flavescentes apice attenuatae non incrassatae, setae exteriores in seriebus indistinctis brevioribus 0.3-0.5 mm longae. Grana pollinis 30-37 μ diam.

TYPE: COSTA RICA: Cartago: steep banks above the Río Grande de Orosi about 9 km southeast of the bridge at Tapantí at an elevation of about 1600 m. Shrub 2 m tall. Ray and disk corollas bright yellow. 20 Dec. 1974. R.L. Wilbur and J.L. Luteyn 18548 (Holotype DUKE; Isotypes CR, US). PARATYPE: COSTA RICA: Cartago: Hwy. 224 on property of ICE hydroelectric plant ca. 20-24 km east of the church in Orosi, beyond the town of Tapantí. Steep roadside banks and disturbed sites adjoined by virgin cloud forests. Elev. ca. 1500-1800 m. Shrub to 3-4 ft. tall. Ray florets yellow. Disk florets yellow, abruptly flaring. Pappus bristly, light brown. Anthers brown. 3 Jan. 1974. F. Almeda Jr. 2378 with M. Flowers, R. Promack, J. Utley, K. Utley, and R. Wyatt (DUKE, US).

Munnozia wilburii has the sagittate leaves and pyramidally paniculate inflorescence with long peduncles that indicate relation to the subgroup of Munnozia containing M. senecionidis. Most members of the group of species differ by the more equal length of the involucre bracts or by different leaf width along with other details of floral size and structure. The probable closest relative is M. senecionidis but a number of differences separate the new species including the prominently winged petiole, the less distinct trinervation at the base of the leaf blade, the presence of more alternate branching in the distal parts of the inflorescence, the more strongly differentiated herbaceous tips of the involucre bracts, the puberulous rather than pilose tubes of the disk corollas and the short throats of the disk corollas. The pubescence of the achene is also comparatively sparse.



PLANTS OF CENTRAL AMERICA

COLLECTED FOR THE UNIVERSITY OF
DURHAM UNIVERSITY

MUNNOZIA WILBURII H. Robinson HOLOTYPE

Costa Rica: Provincia de Cartago: steep
banks above the Río Grande de Orosi about
9 km. southeast of the bridge at Tapantí
at an elevation of about 1600 m.

Shrub 2 m. tall. Ray and disk corollas
bright yellow.

S. L. WILSON #18548
James L. Luteyn

20 December 1974

Munnozia wilburii H. Robinson, Holotype, Duke
University. Photo by Victor E. Krantz, Staff
Photographer, National Museum of Natural History.



Robinson, Harold Ernest. 1978. "Studies in the Liabeae (Asteraceae). XII. A new species of Munnozia from Costa Rica." *Phytologia* 39, 331–333.

<https://doi.org/10.5962/bhl.part.7620>.

View This Item Online: <https://www.biodiversitylibrary.org/item/47392>

DOI: <https://doi.org/10.5962/bhl.part.7620>

Permalink: <https://www.biodiversitylibrary.org/partpdf/7620>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.