

NEED FOR FURTHER STUDY

Because of the rapid expansion of *Orconectes virilis* in much of Utah, expansion of this species must be closely monitored and its impacts upon sportfisheries and native fauna should be documented. In addition to its potential impact on native crayfish, there are indications of negative effects of dense populations with rainbow trout recruitment and early growth (Hepworth and Duffield, in press), and near elimination of aquatic vegetation (Dean 1969). The effects of extensive removal of vegetation on invertebrate production and availability and diversity of littoral zone fishery habitats also require study.

The rash of surreptitious stockings of crayfish in recent years demonstrates an obvious need for a thorough public information program regarding the possible consequences of indiscriminate introductions. Such a program, to be most effective, requires basis in fact and would alone justify well-conceived research into crayfish population dynamics and ecosystem interactions.

Crayfish distribution is especially poorly documented in the western basins of Utah. Furthermore, the collection of *Pacifastacus leniusculus* in Utah County raises the question of whether that species is more widely distributed around Utah Lake or in the western basins. Any crayfish specimens from Utah County or the western basins would therefore be of particular interest.

ACKNOWLEDGMENTS

I am grateful to Dr. Horton H. Hobbs, Jr., U.S. National Museum of Natural History, Smithsonian Institution, Washington, D.C., for confirming the identity of specimens from my reference collection, for identifying *Procambarus clarkii*, and for his helpful comments. Specimens were collected by virtually every regional or research UDWR biologist. In addition, specimens and collection records from the Colorado River drainage were provided by V. Lamarra and R. Valdez of Ecosystem Research Institute, Logan, Utah; R. Radant, Nongame Section, UDWR, and H. Tyus, USFWS, Vernal, Utah.

This work was accomplished in conjunction with fishery studies funded by Federal Aid to Fish Restoration Projects F-43-R and F-44-R.

LITERATURE CITED

- BOUCHARD, R. W. 1976. Geography and ecology of crayfishes of the Cumberland Plateau and Cumberland Mountains, Kentucky, Virginia, Tennessee, Georgia, and Alabama. Part I. The genera of *Procambarus* and *Orconectes*. *Freshwater Crayfish* 2: 563-584.
- . 1977. Distribution, systematic status and ecological notes on five poorly known species of crayfishes in western North America (Decapoda: Astacidae and Cambaridae). *Freshwater Crayfish* 3: 409-423.
- DEAN, J. L. 1969. Biology of the crayfish *Orconectes causeyi* and its use for control of aquatic weeds in trout lakes. U.S. Bureau of Sportfisheries and Wildlife, Technical Paper 24. 15 pp.
- ENG, L. L., AND R. A. DANIELS. 1982. Life history, distribution, and status of *Pacifastacus fortis* (Decapoda: Astacidae). *California Fish and Game* 68(4): 197-212.
- HEPWORTH, D. AND D. DUFFIELD. In press. Impact of an exotic crayfish on stocked rainbow trout in Newcastle Reservoir, Utah. *North American Journal of Fisheries Management*.
- HOBBS, H. H., JR. 1976. Crayfishes (Astacidae) of North and Middle America. U.S. Environmental Protection Agency Water Pollution Control Research Series, 18050 E1D05/72. Cincinnati, Ohio. 173 pp.
- JONES, P. D., AND W. T. MOMOT. 1981. Crayfish production, allochthony, and basin morphometry. *Canadian Jour. Fish. and Aqu. Sci.* 38: 175-183.
- MAY, B., C. THOMPSON, AND S. GLOSS. 1975. Impact of threadfin shad (*Dorsoma petenense*) introduction on food habits of four centrarchids. *Utah Division of Wildlife Resources, Publ.* 75-4. 22 pp.
- PENNAK, R. W. 1978. *Freshwater invertebrates of the United States*. John Wiley and Sons, New York. 803 pp.
- PETTENGILL, T., S. BRAYTON, AND J. JOHNSON. 1984 annual report. Flaming Gorge fishery investigations. *Utah Division of Wildlife Resources, Publ.* 84-07. 55 pp.
- SAKAGUCHI, D. K. 1984. Considerations for managing with crayfish in the waters of the Central Region. *Utah Division of Wildlife Resources, Central Regional Office, Springville, Utah. Memo report.* 6 pp.
- SCHWARTZ, F. J., R. RUBELMANN, AND J. ALLISON. 1963. Ecological population expansion of the introduced crayfish, *Orconectes virilis*. *Ohio Jour. of Sci.* 63(6): 266-273.
- UNGER, P. A. 1978. The crayfishes (Crustacea: Cambaridae) of Colorado. *Natural history inventory of Colorado No. 3.*, University of Colorado Museum. 20 pp.

CRYPTOGAMIC SOIL CRUSTS: RECOVERY FROM GRAZING NEAR CAMP FLOYD STATE PARK, UTAH, USA

Jeffrey R. Johansen¹ and Larry L. St. Clair²

ABSTRACT.—The effects of grazing on the cryptogamic and vascular plant communities at two sites near Camp Floyd State Park, Utah County, Utah, were studied. The grazed site was subject to heavy grazing up until seven years prior to the study. The ungrazed site within the park boundaries had been protected from grazing for 20 years prior to the study and had a well-developed algal-lichen-moss crust. We found that the algae of the grazed site had recovered in terms of degree of crusting. There were no significant differences in the algal communities of the two sites when prevalent species were used as blocks in the ANOVAR analysis. However, when major algal groups were used as blocks, the analysis was significant, with the more recently grazed site having lower algal frequency. This difference, together with a few compositional differences, indicates that, although the algal community seven years following grazing is very similar to the algal community free of grazing for 20 years, the seven-year site is still in the process of recovery and community development. The diatom collections had a higher density in the grazed site, though the difference was not significant. Recovery of the lichen and moss community was not complete. The lichen *Collema tenax* and the mosses *Bryum pallescens* and *Tortula ruralis* were all significantly more abundant in the ungrazed area. Total cover of the lichen and moss components of the soil crusts was significantly lower in the more recently grazed area. Vascular cover was also lower.

Cryptogamic soil crusts are an important component of many arid rangeland ecosystems in the western United States. Such crusts have been found to be important in nitrogen fixation (Snyder and Wullstein 1973, Rychert and Skujins 1974) and enhancement of seedling establishment (St. Clair et al. 1984). The greatest benefit of cryptogamic crusts, however, is probably reduction of soil erosion. Soil aggregation, particularly by blue-green algae, reduces detachment of soil particles by wind and rain (Bailey et al. 1973, Anantani and Marathe 1974). Improved water penetration in crusted soils reduces runoff and subsequent erosion (Brotherson and Rushforth 1983). Sedimentation is also reduced by the increased tortuosity of surface water pathways due to the characteristic hummocking of desert crusts.

The effects of both burning and grazing on soil cryptogams have recently been investigated. Range fires can severely damage all components of the soil crust (Johansen, Javakul, and Rushforth 1982, Johansen et al. 1984, Callison et al. 1985). Several workers have noted that moderate to heavy grazing can seriously damage soil cryptogams because the crusts are trampled by livestock. Rogers and

Lange (1971) noted that stocking pressure was negatively correlated with lichen cover and that soil mobility and erosion was increased in areas of reduced lichen cover. Kleiner and Harper (1972) found that the effects of grazing were more notable on the soil cryptogams than on the vascular plant communities in Canyonlands National Park. They also noted that soil erosion was higher in areas with lower cryptogam cover and observed changes in organic matter, available phosphorus, and calcium in eroded soils (Kleiner and Harper 1977). The effects of long-term moderate to heavy grazing near Navajo National Monument also showed that grazing has a more pronounced effect on the cryptogamic cover and diversity than on vascular plant cover and diversity (Brotherson et al. 1983). Burros in the Grand Canyon are currently causing erosion through destruction of *Tortula* (bryophyte) and lichen crusts (Phillips et al. 1977).

Several factors influence the development of cryptogamic crusts. Cryptogamic growth is best in soils of high electrical conductivity, high phosphorus, and high silt content (Anderson et al. *Factors*, 1982). Crust buildup is also positively correlated with soil alkalinity. The crusts of gypsiferous soils of southern

¹SERI, 1617 Cole Blvd., Golden, Colorado 80401.

²Department of Botany and Range Science, Brigham Young University, Provo, Utah 84602.

Utah are particularly well developed, having high lichen and algal diversity (Anderson and Rushforth 1976). Recovery of crusts from disturbance follows several trends. Algae are the most resistant component to disturbance (Anderson et al. *Recovery*, 1982) and are also the quickest to recover (Johansen et al. 1984). Lichens and mosses are slower to recover. Anderson et al. *Recovery*, (1982) found that lichens and mosses had become fairly well established after a 14–17-year period of protection from grazing. Exclosures protected for 37–38 years showed little change in the moss flora, though lichen diversity was greater than in the 14–17-year-old exclosures. This paper also indicated a need for studies examining recovery in the first 15 years following protection from disturbance.

The primary purpose of our study was to determine the degree of recovery of the cryptogamic crust community in a soil that has been protected from grazing for seven years near Camp Floyd State Park. This study differs from past studies in that the grazed area is compared with an adjacent area that has been protected for 20 years rather than with an area still under grazing pressure.

SITE DESCRIPTION

The study was conducted near and within the boundaries of Camp Floyd State Park, Utah County, Utah. The park is located 0.4 km southwest of Fairfield, Utah, along State Highway 73 in central Utah and is relatively small (16.2 hectares). It was established in 1962 with the principal point of interest being an inactive military cemetery which occupies only a small portion of the total area of the park. The balance of the park has not been developed and is dominated by an *Atriplex confertifolia*–*Sarcobatus vermiculatus* desert shrub community. A well-developed cryptogamic soil crust flora consisting of various species of lichens, mosses, and algae is present. The soils at Camp Floyd belong to the Woodrow silt loam with an average water-holding capacity of 28–30 cm of water for the 1.5 m (5 ft) profile. This soil type has a slow permeability and is classified as a mixed (calcareous) mesic, xeric torifluent. The mean annual rainfall in this area is 35 cm.

From 1935 until the establishment and subsequent fencing of the park area, the 1,600

hectares of shrubland immediately around the cemetery were heavily grazed by sheep and cattle during winter months (October to May). Fencing of the park property has effectively eliminated grazing in the park since 1962. This has permitted the establishment of a diverse and well-developed algal-lichen-moss community. The area outside the park continued to be heavily grazed until 1975, at which time livestock were removed from this range.

METHODS

Two permanent transects with points placed every 2 m were established in the vicinity of Camp Floyd State Park. The ungrazed area transect consisted of 6 points within the park boundaries. Data from this transect were compared with similar data collected from burned sites, and that comparison is the basis of a previous paper (Johansen et al., 1984). The grazed area transect consisted of 8 points one mile south of the park. Fieldwork was conducted from September to November, 1982.

Two crust samples were collected from each transect point at opposite compass points 0.25 m from the center of the point on 23 September 1982. Each sample was prepared for culturing by gently crushing soil clods to a maximum diameter of 5 mm. Twenty cm³ of soil from each sample were placed in a sterilized petri dish and saturated with 20 ml of distilled water. Samples were then incubated under continuous cool-white fluorescent light at room temperature (24 C) for 10 days. Percent visible algal cover in each petri dish was estimated at the end of the culture period. Frequency and relative abundance of living algal species were also estimated at that time by subsampling the center of each petri dish and examining this subsample under the light microscope. A total of 25 microscope fields were examined for each subsample, and presence and absence of each species in each field were noted. Permanent diatom mounts were prepared using standard acid oxidation procedures and Hyrax diatom mountant. Slides were prepared using quantitative dilutions for quantitative comparisons of soil diatoms (Johansen et al. 1982).

Visual estimates of total cryptogamic and vascular plant cover were made in the field

TABLE 1. Average percent frequencies of living algal species in grazed and ungrazed areas. Each species was tested using the Mann-Whitney test to determine if significant differences existed between treatments (* indicates $p < .05$, ** indicates $p < .01$). Estimated visible algal cover in culture dishes is also given.

Species	Grazed	Ungrazed
CYANOPHYTA		
<i>Anabaena variabilis</i> Kuetzing	.3	4.0
<i>Aphanothece castagnei</i> (Breb.) Rabh.	1.0	.3
<i>Chroococcus minor</i> (Kuetz.) Naegeli	1.0	.3
<i>Chroococcus turgidus</i> (Kuetz.) Naegeli		2.0
<i>Gloeotheca linearis</i> var. <i>composita</i> G. M. Smith		.3
<i>Microcoleus vaginatus</i> (Vauch.) Gomont	55.0	57.0
<i>Nostoc commune</i> Vaucher	.5	10.0 *
<i>Nostoc muscorum</i> C. A. Agardh	.3	
<i>Nostoc</i> cf. <i>paludosum</i> Kuetzing	.8	1.7
<i>Nostoc</i> species	5.5	7.7
<i>Phormidium minnesotense</i> (Tild.) Drouet	27.8	15.3
<i>Synechococcus aeruginosus</i> Naegeli	.3	
<i>Tolypothrix tenuis</i> (Kuetz.) Schmidt	1.8	3.7
Unknown Chroococcaceae	1.5	4.0
CHLOROPHYTA		
<i>Ulothrix tenerrima</i> Kuetzing	.3	3.0
Unknown coccoids	16.3	23.0
BACILLARIOPHYTA		
<i>Hantzschia amphioxys</i> (Ehr.) Grunow	18.5	26.0
<i>Navicula mutica</i> Kuetzing	5.0	12.3
<i>Navicula paramutica</i> Bock	1.0	.3
<i>Pinnularia borealis</i> Ehr.	.8	.7
FLAGELLATES	.3	1.7
TOTAL SUM FREQUENCY	137.0	173.3
VISIBLE ALGAL COVER	16.6	36.5 **
SHANNON-WIENER DIVERSITY INDEX	2.35	2.75**

using a 1/4 m² circular quadrat placed at each permanent transect point. In addition to the 1/4 m² circular quadrats, a smaller 2 × 50 cm rectangular quadrat consisting of ten 2 × 5 cm subquadrats was placed perpendicular to the transect at each point and percent cover of algal crust and lichen and moss species was estimated. Cover was measured in November in the ungrazed area and in September in the more recently grazed area.

Soils were analyzed by the Soil Analysis Laboratory, Department of Agronomy and Horticulture, Brigham Young University, for pH, phosphorus, nitrate, Kjeldahl nitrogen, potassium, calcium, magnesium, zinc, iron, manganese, sodium, copper, organic matter, electrical conductivity, and sodium adsorption ratio. All samples were taken from the top 17.5 cm of soil for these analyses.

Mann-Whitney tests (Ryan et al. 1976) were performed for each species to determine significance levels of differences between grazed and ungrazed areas. This test was used in preference to the Students t-test (Snedecor

and Cochran 1980) because some of our data were not normal. Shannon-Wiener diversity indices were calculated for the living algae and diatom communities for each transect point (Patten 1962, Shannon and Weaver 1949). Similarity indices for all 14 stands were calculated following the methods of Ruzicka (1958). Four different sets of similarity indices were calculated using lichen and moss cover, living algal density, diatom density, and vascular plant cover. These four sets of indices were then clustered following Sneath and Sokal (1973) to illustrate the degree of similarity between grazed and ungrazed stands in the four different plant communities. Importance values for species of living algae and diatoms were determined by multiplying average percent relative density times presence (Warner and Harper 1972). Species with importance values above 1.00 were used in the subsequent ANOVAR analyses.

Multivariate analysis of variance adapted in preference to the Students t-test (Snedecor for a fixed effects, unbalanced design follow-

ing the methods of Bryce et al. (1980) was used to analyze differences between blocks (lichens, mosses, algae, vascular plants) and treatments (time since grazing). Several separate ANOVAR tests were run, including tests using the vascular plant cover data, lichen and moss cover data, living algal data, and subfossil diatom data. In some data sets the variance of each species and group was related to the mean. To satisfy the homogeneity of variance assumption of analysis of variance for these data sets, a $\log(x+1)$ transformation was used (Bartlett 1947). With each analysis of variance, the standardized residuals were plotted against the normal scores to give a measure of normality. In all cases the probability plot thus generated was subjectively judged as being normal or close to normal. Our use of transects satisfied the requirements of systematic sampling as described by Cochran (1977). The Duncan multiple range test was used to determine significance of differences between species means when analysis of variance showed significance for this factor (Duncan 1955). Unless otherwise stated, the alpha level used for this test was 0.01.

RESULTS

Living Algae

The seven most important living algal species were *Microcoleus vaginatus* (importance value=55.86), *Phormidium minnesotense* (20.83), unknown green coccoid (18.46), *Hantzschia amphioxys* (17.06), *Nostoc* species (4.59), *Navicula mutica* (4.07), and *Nostoc commune* (1.42). Average percent frequencies for all algal species are given in Table 1. Multivariate analysis of variance based on the above seven prevalent species showed that the algal communities of the two areas were not significantly different. The Mann-Whitney test on the visual algal cover estimated in the field supported this conclusion (Table 3). The multivariate analysis did show that the species had significantly different frequencies. Duncan's test showed that *Microcoleus vaginatus* was significantly more abundant than all other algal species. *Phormidium minnesotense*, unknown coccoid green algae, and *Hantzschia amphioxys* were significantly more abundant than all less common algal species. The interaction between treatment and species was significant ($p=.023$). This was likely due to the




Fig. 1. Interaction between living algal species and treatment, significant at $p=.023$, using multivariate analysis of variance.

fact that *Hantzschia amphioxys*, *Navicula mutica*, *Nostoc commune*, and coccoid green algae were all more abundant in the park area, whereas *Phormidium minnesotense* was more numerous in the more recently grazed stands (Fig. 1). *Microcoleus vaginatus* and *Nostoc* species showed only minor differences in frequency between the two sites.

A separate analysis of variance was conducted to compare major algal groups. When using this data set, the difference between treatments was significant ($p=.041$), the more recently grazed plots having lower values. The groups were also significantly different ($p<.001$). Duncan's test showed that Cyanophyta were significantly more abundant than the other three algal groups. Bacillariophyceae and Chlorophyta were significantly more abundant than unidentified flagellates.

Nostoc commune had a significantly greater population in the 20-years-since-grazing area ($p=.024$) according to the Mann-Whitney test. No other living algal species or groups were significantly different at the two sites according to this test.

Subfossil Diatoms

The nine most important subfossil diatom species, including chrysophyte cysts observed in diatom mounts, were *Hantzschia amphioxys* (importance value 69.93), *Navicula mutica* (54.96), chrysophyte cysts (46.43), *Pinnularia borealis* (18.91), *Navicula mutica* var. *cohnii* (6.06), *Navicula paramutica* (3.65), *Navicula mutica* var. *nivalis* (2.42), *Navicula contenta* f. *parallela* (2.39), and *Cy-*



1986. "New species and new records of North American Pityophthorus (Coleoptera: Scolytidae). Part 6. The lautus group." *The Great Basin naturalist* 46, 641–645. <https://doi.org/10.5962/bhl.part.7740>.

View This Item Online: <https://www.biodiversitylibrary.org/item/33077>

DOI: <https://doi.org/10.5962/bhl.part.7740>

Permalink: <https://www.biodiversitylibrary.org/partpdf/7740>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Brigham Young University

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.