

If the hollow interior be the natural condition of the *Mylodon*-like humerus under inspection, it would not belong to *Mylodon robustus*. Independently of the cavity indicated, the bone is sufficiently different in size and form to indicate a different species from the *Mylodon Harlani* of North America. The humerus from Oregon, described by Perkins (Am. Jour. Sci. 1841, xlii, 136), and referred to the latter by Prof. Owen, is not only much larger, but it is of greater breadth in relation with its antero-posterior diameter. The fragment of a humerus from Big-Bone-Lick, Ky., represented in fig. 3, plate xiv of my "Memoir on the Extinct Sloth Tribe," is somewhat smaller than the corresponding part of the Oregon specimen, and is more compressed or wider in comparison with the antero-posterior diameter.

Prof. Leidy further observed that there appeared to be a point of some significance in the anatomy of the mandible of *Dromatherium silvestre* worthy of attention, though the appearance may turn out to be a deceptive one. Prof. Emmons had discovered three isolated rami of mandibles of this most ancient of American mammals in the triassic coal of North Carolina. Of the specimens, one is represented in fig. 66 of Emmons' American Geology, repeated in outline in fig. 650 of Dana's Geology. Another specimen Prof. Emmons presented to the Academy, and is contained in our museum. The point of interest to which reference is made is the apparent absence of a condyle. This process may have been lost, but in the two specimens seen by Prof. L.—that figured by Prof. Emmons, and that preserved in our museum—a separation of the process is not obvious.

March 8th.

DR. CARSON, Vice-President, in the Chair.

Twenty-five members present.

PROF. LEIDY made the following remarks:—The reptilian remains from the cretaceous formation near Fort Wallace, Kansas, presented to the Academy by Dr. T. H. Turner, and described by Prof. Cope under the name of *Elasmosaurus platyurus*, belong to an Enaliosaurian, as originally suggested by Prof. Cope. The anatomical characters of the different regions of the vertebral column, those of the shoulder and pelvic girdles, and of the preserved portions of the skull and teeth, are decidedly Plesiosaurian.

Prof. Cope has described the skeleton in a reversed position to the true one, and in that view has represented it in a restored condition in fig. 1, pl. ii. of his "Synopsis of the Extinct Batrachia and Reptilia," Pt. I, August, 1869, published in advance for the fourteenth volume of the Transactions of the American Philosophical Society. To explain the apparently anomalous and reversed arrangement of the articular processes (zygapophyses) of the vertebræ, he has supposed that those as ordinarily existing are substituted by the second set of articular processes (zygophene and zygantrum), as found in serpents and iguanians (Proc. Bost. Nat. Hist. Soc. xii, 265; Syn. Ext. Bat. and Rept. 42).

The finding of a portion of the jaws, as reported by Dr. Turner, in the vicinity of what Prof. Cope has supposed to be the cervical portion of the skeleton, and which he considers as confirmatory of the view he has taken of its position, without further consideration, is more than compensated in the opposite end of the column terminating in a coössified axis and atlas, as is the case also in the mature *Plesiosaurus*. The cup of the atlas still retains the hemispherical occipital condyle.

The Kansas saurian was wonderful for the length of its neck, far exceeding in this respect the *Plesiosaurus*. The vertebræ in the specimen form a nearly unbroken series to the seventy sixth inclusive. If we regard all as cervical until the transverse processes begin to spring in part from the spinal arch, it 1870.]

will comprise the extraordinary number of seventy-two. In the different species of *Plesiosaurus*, so far as known, the number ranges from twenty-four to forty-one. The length of the neck, independent of the head, was about twenty-two feet.

The cervical vertebræ successively increase in length to about the forty-fourth, then remain nearly the same to the sixtieth, and afterwards gradually decrease. The atlo-axis is about $2\frac{1}{4}$ inches long; the third cervical is $1\frac{1}{2}$ inches; the tenth nearly 2 inches; the twentieth $2\frac{3}{4}$ inches; the thirtieth $3\frac{1}{2}$ inches; the fortieth 4 inches; the forty-fourth $4\frac{1}{2}$ inches, and so to the sixtieth; and the sixty-eighth to the last one about $3\frac{3}{4}$ inches, which is also about the length of the succeeding four dorsals.

The imperfections in the remainder of the vertebral column of the Kansas saurian do not permit a positive estimate to be made of the comparative extent of the trunk and tail.

A comparison of the caudal vertebræ with isolated specimens from the cretaceous formations of Alabama, Mississippi and New Jersey, leaves but little doubt that *Elasmosaurus* is identical with *Discosaurus*. Such also appears originally to have been the opinion held by Prof. Cope in regard to a portion of the same skeleton, which he referred to a species with the name of *Discosaurus carinatus* (LeConte's Notes on the Geology of the Survey of the Union Pacific Railroad, 1868, p. 68).

Specimens of vertebral bodies from the New Jersey green sand, referred to *Cimoliasaurus* (Cret. Rept. of the United States, pls. v, vi), and supposed by me to belong to the posterior part of the column, are seen by comparison with the Kansas skeleton to be cervical and perhaps anterior dorsals. The difference in the proportions of the corresponding vertebræ appear to indicate the genus to be distinct from *Discosaurus*.

The imperfect vertebral specimens from Arkansas, originally referred to *Brimosaurus* (Pr. Acad. Nat. Sci. 1854, 72, pl. ii, figs. 1—3), are probably posterior cervicals of *Discosaurus*.

In the true view of *Discosaurus* and its allies, the so-called order of *Streptosauria* (Proc. Bost. Nat. Hist. Soc. 1869, 265; Synopsis Ext. Batr. and Rept., 40) fails to maintain its position.

The extensive shoulder and pelvic girdles of the Kansas saurian, so much like those of *Plesiosaurus*, were most probably provided with limbs constructed like those of the latter animal.

In its restored condition *Discosaurus* would appear to have resembled *Plesiosaurus* in its form as ordinarily represented, excepting that it possessed a much longer neck,—one indeed that exceeded that of all known animals. We may imagine this extraordinary creature, with its turtle-like body, paddling about, at one moment darting its head a distance of upwards of twenty feet into the depths of the sea after its fish prey, at another into the air after some feathered or other winged reptile, or perhaps, when near shore, even reaching so far as to seize by the throat some biped dinosaur.

Prof. Leidy subsequently exhibited fragments of a fossil mandible, upon which he made the following remarks: The specimens were obtained by Prof. F. V. Hayden from the tertiary rocks of the Bridger Group, near Fort Bridger, Wyoming. They indicate a carnivorous animal, apparently of an extinct and heretofore unknown genus. Unfortunately the teeth are lost, except portions of some of the molars. The animal was larger than our Panther, and probably was a member of the same family, but with a relationship to the Hyænas. The portion of the jaw preserved nearly agrees in form with the corresponding portion in the Cats, but its depth below the position of the teeth is proportionately greater, resembling in this respect more the condition in the Striped Hyæna.

Remains of the canine alveolus indicate a tooth of more robust proportions than in the latter animal. The symphysis pursues the course of the alveolus.

[March,

Of several mental foramina, the largest one is situated below the position of the first premolar.

The condyle holds the same relative height as in Cats. The back portion of the jaw, including the coronoid process, is proportionately not so broad as in the latter. The masseteric fossa is shallower at its lower part, and is not bounded by the everted base, but is defined a little above the latter in a curvature sweeping from the condyle downward and forward to an angle about half-way below the position of the last molar tooth.

The molar teeth, of which there were five, completely occupied the space back of the canine, as in some of the viverrine and musteline animals, there being no hiatus in the series. All the molars were double-fanged, and none appear to have been of the tubercular kind.

The first premolar was the smallest, and the third premolar appears to have been the largest tooth of the molar series. The fourth premolar was intermediate in size to the third and the last molar, which appears to have but little exceeded this and the second premolar.

The remains of the crown of the last molar indicate a bilobed tooth, apparently like the sectorial molar of *Felis*, and without a heel. The crown of the tooth in advance was provided with a well-developed heel, but the fore part is too much broken to ascertain its form. The larger tooth in advance, the third premolar, retains its heel, which has a subtrenchant fore and aft border, and is bounded internally and externally by an oblique basal ridge.

For the animal indicated by the fossil jaw, the name of *Patriofelis ulta* is proposed. The measurements of the specimen are as follow:

Estimated length of lower jaw when complete	6 inches.
Breadth of coronoid process at base	$1\frac{3}{4}$ "
Height at condyle, and below last molar tooth.....	$1\frac{3}{8}$ "
Height below first premolar.....	$1\frac{1}{2}$ "
Length of molar series.....	3 "

Breadth of crown of first molar tooth, $4\frac{1}{2}$ lines; second do., 8 lines; third do., 9 lines; fourth do., $7\frac{3}{4}$ lines; fifth do., 8 lines.

PROF. MARSH exhibited specimens of the remains of a bird allied to the Turkey, which he considered as belonging to an extinct species. These remains were said to have been found in the green sand of Monmouth Co., N. J., but doubtless were from the post-tertiary above it. He proposed to name the species *Meleagris altus*.

Prof. Marsh also called attention to the tooth of a Peccary from the miocene of Shark River, New Jersey. It is a second molar from the left side of the lower jaw. He proposes to call the species to which it belonged *Dicotyles antiquus*.

March 15th.

The President, DR. RUSCHENBERGER, in the Chair.

Twenty members present.

The following paper was presented for publication:

"Cross Fertilization and Law of Sex in *Euphorbia*." By Thomas Meehan.

March 22d.

DR. CARSON, Vice-President, in the Chair.

Thirty-six members present.

The following paper was presented for publication:

1870.]



1870. "March 8th." *Proceedings of the Academy of Natural Sciences of Philadelphia* 22, 9–11.

View This Item Online: <https://www.biodiversitylibrary.org/item/18247>

Permalink: <https://www.biodiversitylibrary.org/partpdf/85457>

Holding Institution

MBLWHOI Library

Sponsored by

MBLWHOI Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.