

the fifth egg was laid a dog came around near the back door and caught sight of the bird in the shoe and made a dash for her, the bird escaping, but breaking three of the eggs.

The shoe, nest, and two eggs are in my collection. The nest was composed of dry grasses and fine moss and lined with horse hair. — ARTHUR WILLIAM BROCKWAY, *Old Lyme, Conn.*

**Puerto Rico Honey Creeper.** — I have been interested in a pair of Honey Creepers, *Cæreba portoricensis* (Bryant), building about my house. They began in a rose bush, but it being too close to the ground they deserted the place and are now busy upon another nest in a small tree. The nest is a little larger than a baseball, perfectly round, with the opening like a well drilled auger hole, just below the middle. Outside are grasses and bits of twine; inside are feathers, and when the birds leave the nest for any time they cover the hole with a couple of feathers. The female does this also when she is within, just peeping out with bill and head, which with the aid of my glass makes a *real picture*. These birds are our 'Jenny Wrens,' and there are a good many of them here all around our houses, especially where roses, coral plants, and other smaller flowers abound. We have become very much attached to them. — GEORGE B. PRATT, *San Juan, Puerto Rico.*

**Notes on Marian's Marsh Wren, *Cistothorus marianæ*, and Worthington's Marsh Wren, *Cistothorus palustris griseus*.** — On April 16, 1897, I shot a very dark colored Marsh Wren near Mount Pleasant, South Carolina, which has the top of head deep black. At Mr. Ridgway's request, I sent the specimen to him and under date of June 10, 1897, he wrote me as follows: "I have just finished examining the Wren, with the aid of Mr. Brewster's series of both *marianæ* and *griseus*. Your dark bird is intermediate, but on the whole may best be considered an exceptionally dark specimen of *griseus*. *Marianæ* is a more rusty brown and usually has the breast speckled with blackish."

I sent the Wren above mentioned to Mr. Brewster and he wrote me under date of Dec. 6, 1897, as follows: "As to the April bird (Wren), I cannot agree with Mr. Ridgway, for it has absolutely none of the true characters of *griseus* and is quite indistinguishable, so far as I can see, from some of my examples of *marianæ* from Tarpon Springs, Florida; although it certainly has more white beneath than is usually the case with that form. In respect to the coloring of the upper parts, and especially of the crown, it is typical *marianæ*, to which I should accordingly refer it despite the locality at which it was taken."

It will be seen from the above that this record extends the range of this bird to the Atlantic coast, and that it is an abundant bird in this State during the migrations the following records will show. On Oct. 1, 1898, I killed four; Oct. 4, five; Oct. 11, one; Oct. 15, one; Oct. 17, three; Oct. 28, two; Oct. 31, two; April 21, 1899, one; May 6, one; May 8, one.



1899. "Puerto Rico Honey Creeper." *The Auk* 16, 361–361.

<https://doi.org/10.2307/4069375>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/54100>

**DOI:** <https://doi.org/10.2307/4069375>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/89011>

#### **Holding Institution**

Smithsonian Libraries and Archives

#### **Sponsored by**

Smithsonian

#### **Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.