

# New Records for Western Pacific *Morum* (Gastropoda: Harpidae) with Biogeographic Implications

by

WILLIAM K. EMERSON

Department of Invertebrates, American Museum of Natural History,  
New York, New York 10024, USA

**Abstract.** Major extensions of the known range are reported for six species of the prosobranch genus *Morum*, namely: *M. teramachii*, *M. uchiyamai* and *M. joelgreeni* in the Mariana Islands, *M. uchiyamai* and *M. bruuni* in the region of New Caledonia, *M. cancellatum* in the Fiji Islands, and *M. kurzi* in the Solomon Islands. The distributional patterns of the 15 recognized species of *Morum* living in the Indo-West Pacific biogeographic region are evaluated in terms of the occurrences of these taxa on the regional lithospheric plates. The fossil and modern distributional patterns of *Morum* (*sensu lato*) suggest that these gastropods are remnants of a Tethyan faunal element which is limited in distribution owing largely to the apparent lack of teleplanic larvae.

## INTRODUCTION

This paper records major extensions of the known range for six Indo-Pacific species of the genus *Morum*. Specimens of *Morum teramachii* Kuroda & Habe, 1961, *M. uchiyamai* Kuroda & Habe, 1961, and *M. joelgreeni* Emerson, 1981, were obtained in the Mariana Islands by deep-water shrimp trapping operations of the NOAA vessel *Townsend Cromwell* during 1982 to 1984. This survey was conducted by the National Marine Fisheries Service, Southwest Fisheries Center Honolulu Laboratory as part of their Resource Assessment Investigation of the Mariana Archipelago Program (EMERSON & MOFFITT, 1988). Specimens of *Morum bruuni* (Powell, 1958) and *M. uchiyamai* Kuroda & Habe were dredged in 1986 off New Caledonia in deep water by the N.O. *Vauban* and N.O. *Coriolis*, vessels operated by the ORSTOM Center in Nouméa, New Caledonia (BOUCHET, 1986; RICHER DE FORGES, 1988). A single specimen of *M. kurzi* Petuch, 1979, was collected in 1985 by SCUBA diving from off Guadalcanal in the Solomon Islands. A specimen of *M. cancellatum* B. G. Sowerby I, 1824, inhabited by a hermit crab was taken in a baited trap off Suva Reef, Fiji Islands, in 1979.

These new records significantly extend the known ranges of these taxa eastward along the Eurasian, Philippine, and

Indian-Australian Lithospheric Plates to the western margin of the Pacific Plate.

## ABBREVIATIONS

The following abbreviations for institutions and organizations are used in the text.

AMNH—American Museum of Natural History, New York

BM(NH)—British Museum (Natural History), London  
MNHN-Paris—Muséum National d'Histoire Naturelle, Paris

NMFSHL—National Marine Fisheries Service Southwest Fisheries Center, Honolulu Laboratory

NMNH—National Museum of Natural History, U.S. National Museum collection (USNM), Smithsonian Institution, Washington, D.C.

ORSTOM—Institut Français de Recherches pour le Développement en Coopération, Paris and Nouméa

TC—National Oceanic and Atmospheric Administration vessel *Townsend Cromwell*

UGMLM—University of Guam, Marine Laboratory, Mangilao, Guam

USBF—United States Bureau of Fisheries






Figure 1

Distribution of *Morum* on the Pacific Plate (species numbers 6–8) and on or near the border of the Pacific Plate (species numbers 1–5, 7). 1 = *M. teramachii*, 2 = *M. uchiyamai*, 3 = *M. joelgreenei*, 4 = *M. bruuni*, 5 = *M. cancellatum*, 6 = *M. kurzi*, 7 = *M. ponderosum*, and 8 = *M. macdonaldi*. (Base map, without the plate boundaries, courtesy of R. T. Abbott.)

## SYSTEMATIC TREATMENT

Family HARPIDAE Bronn, 1849

Subfamily MORUMINAE Hughes & Emerson, 1987

Genus *Morum* Roeding, 1798

Subgenus *Oniscidia* Moersch, 1852

*Morum* (*O.*) *cancellatum* G. B. Sowerby I, 1824

(Figures 6, 7)

**New record:** FIJI ISLANDS: off Suva Reef (18.05°S, 178.25°E), Viti Levu in 220 m (120 fms), 21 March 1979, in baited *Nautilus* trap, 1 crabbed specimen, Roper and Sweeney collectors (USNM 773946).

**Remarks:** This record is based on a single, well-preserved, hermit-crabbed specimen, with fully developed apertural dentition and parietal shield (Figures 6, 7). The present specimen (44.1 mm × 27.5 mm) is slightly smaller than the lectotype, which measures 47.2 mm in height and 28.6 mm in width [BM(NH) 197744, EMERSON, 1985:54, figs. 17, 18], but is otherwise typical of this species.

**Range:** Previously known from southern Japan, Taiwan, the south China coast (off Pratas Island, in 161 m [88 fms], USBF Sta. 5111 (USNM 232801, USNM 237710), and the central Philippine Islands (Sandugan Pt., Signjos, Mindoro Island, AMNH 207315). Here recorded from the Fiji Islands in 220 m (dead specimen).

*Morum* (*O.*) *joelgreenei* Emerson, 1981


(Figures 4, 5)

**Synonym:** *M. (O.) celinamarumai* Kosuge, 1981 (see EMERSON, 1985: 52, pl. 1, figs. 9, 10). **Type locality:** "Bohol, Philippines."

**New record:** MARIANA ARCHIPELAGO: off Pagan Island (18°04.8'N, 145°45.2'E), in 201–284 m (110–150 fms), TC 84-02, Sta. 3, 20 February 1984 (AMNH 232122), from shrimp trap, 1 crabbed specimen.

**Remarks:** The present specimen (Figures 4, 5) is well preserved, but small (measuring 32 mm in length and 20.3





#### Explanation of Figures 2 to 13

Figures 2-13. Specimens representing the six taxa of *Morum* (*Oniscidia*) for which significant range extensions are recorded in this paper. Figures 2, 3: *M. kurzi* (AMNH 213436);  $\times 2$ . Figures 4, 5: *M. joelgreenei* (AMNH 232122);  $\times 1$ . Figures 6, 7: *M. cancellatum* (USNM 773946);  $\times 1$ . Figures 8, 9: *M. uchiyamai* (AMNH 232123);  $\times 1$ . Figures 10, 11: *M. bruuni* (Station DW71, MNHN-Paris);  $\times 1.5$ . Figures 12, 13: *M. teramachii* (AMNH 232124);  $\times 1$ .



mm in width) with an incompletely developed outer lip and an immature parietal shield.

**Range:** Previously known from moderately deep water (100 to 150 m) off the southern Philippine Islands. Here recorded from the Mariana Islands in deeper water (dead specimen).

*Morum (O.) kurzi* Petuch, 1979

(Figures 2, 3)

**New record:** SOLOMON ISLANDS: 1 km west of Honiara, Guadalcanal Island, in 25–30 m (14–17 fms), spring of 1985, 1 specimen, by SCUBA diving, Johnson Kengalu collector (AMNH 213436).

**Remarks:** On a collecting trip to the Solomon Islands, Don Pisor obtained a single, live-collected specimen from a local diver (Figures 2, 3). It is typical of the Philippine specimens (type locality off Panglao, Bohol Island) and is essentially the same size as the holotype (EMERSON, 1985: 53, pl. 1, figs. 5, 6). The present specimen was generously donated to the AMNH by Mr. Pisor.

**Range:** Previously known only from the southern Philippines in depths of 110 (AMNH 213707) to 250 m (PETUCH, 1979). Here recorded from the Solomon Islands in lesser depths (25–30 m).

*Morum (O.) teramachii*

Kuroda & Habe in Habe, 1961

(Figures 12, 13)

**New records:** MARIANA ARCHIPELAGO: off Pagan Island (15°01'N, 145°14'E), in 366 m (220 fms), TC 82-04, Sta. 31, 3 August 1982 (AMNH 232126), from shrimp trap, 1 crabbed specimen; off Anatahan Island (16°21.6'N, 145°43.9'E), in 353 m (193 fms), TC 82-02, Sta. 47, 28 April 1982 (AMNH 232127), from shrimp trap, 2 live-taken specimens; off Farallon de Medinilla (16°08'N, 146°07'E), in 366 m (200 fms), TC 82-03, Sta. 104, 20 June 1982 (AMNH 232128), from shrimp trap, 1 crabbed specimen; off Arakane Reef (15°37.4'N, 142°46.2'E), in 311–476 m (170–260 fms), TC 83-05, Sta. 142, 15 December 1983 (UGMLM, No. 1; AMNH 232129, Nos. 3, 4), from shrimp traps, 4 crabbed specimens, No. 2 (AMNH 232124) here illustrated (Figures 12, 13); off Arakane Reef (15°37.6'N, 142°46.1'E) in 311–476 m (170–260 fms), TC 83-05, Sta. 151, 17 December 1983, USNM 869025, from shrimp trap, 1 crabbed specimen; off Esmeralda Island (15°01'N, 145°14'E), in 357–448 m (195–245 fms), TC 82-03, Sta. 39, 11 June 1982 (AMNH 232130), from shrimp trap, 1 crabbed specimen.

**Remarks:** The shells of the 10 specimens reported here range in measurements from 63.1 mm in length and 34.1 mm in width with 6½ post-nuclear whorls, to 40.9 mm in length and 26.4 mm in width with 5½ post-nuclear whorls.

The figured type specimen of this taxon was stated to measure 55.5 × 32.3 mm and was dredged off Cape Ashiuri, Kochi Pref., Shikoku, Japan, in ca. 200 m (INABA & OYAMA, 1977:121).

**Range:** Previously known from southern Japan and the southern Philippine Islands (GLASS & FOSTER, 1986:68) in moderately deep water (100–200 m). Here recorded from the Mariana Islands, in depths of 170–448 m (live-taken specimens from 353 m).

*Morum (O.) uchiyamai* Kuroda & Habe  
in Habe, 1961

(Figures 8, 9)

**New records:** MARIANA ARCHIPELAGO: off Saipan Island (15°15.83'N, 145°41.10'E), in 366–384 m (200–210 fms), TC 81-01, Sta. 179, 13 April 1981 (AMNH 232123), from shrimp trap, 1 crabbed specimen here illustrated (Figures 8, 9); off Esmeralda Bank (15°01.8'N, 145°13.7'E), in 384–430 m (210–235 fms), TC 84-01, Sta. 16, 8 January 1984 (UGMLM), from pipe dredge, 1 dead and discolored specimen. SOUTH OF NEW CALEDONIA: (22°53'S, 167°11'E) in 375–402 m (206–221 fms), N.O. *Vauban* SMIB2 cruise, Sta. DW15, 18 September 1986, MNHN-Paris, 1 crabbed specimen, Menou and Tirard collectors; (21°01'S, 167°27'E) dredged in 250 m (138 fms), R.V. *Alis* MUSORSTOM 6, Sta. DW453, 20 February 1989, MNHN-Paris, 1 live-collected specimen, Bouchet and Richer de Forges collectors.

**Remarks:** All three of the dead-taken specimens are smaller than the type specimen (51.5 mm × 27.5 mm), which was dredged off Kochi Pref., Shikoku, Japan, in 200–300 m (INABA & OYAMA, 1977:128). The live-taken New Caledonia specimen (52.2 × 28.4 mm) specimen is approximately the size of the holotype. The largest Marianan specimen (45.1 mm × 24.9 mm) is illustrated.

**Range:** Previously known from the region of the East China Sea (southern Japan and Taiwan) in moderately deep water (183–300 m). Here recorded in the Mariana Islands and in the region south of New Caledonia, in depths of 200–250 m (living specimens from 250 m).

*Morum (O.) bruuni* (Powell, 1958)

(Figures 10, 11)

**Synonym:** *Pulchroniscia delecta* Garrard, 1961 (see BEU, 1976: 225–229, figs. 1, 2, 4, 5, 11–15).

**New records:** CORAL SEA: off New Caledonia (22°48'S, 159°24'E), dredged in 360–390 m (198–215 fms), N.O. *Coriolis*, MUSORSTOM 5 Cruise, Sta. 299, 11 October 1986, MNHN-Paris, dead specimen encrusted with barnacles, etc., Bouchet, Metivier and Richer de Forges collectors. SOUTH OF NEW CALEDONIA: (24°42'S, 168°10'E), dredged in 230 m (127 fms), N.O. *Coriolis* CHALCAL 2



Cruise Sta. DW71, 27 October 1986, MNHN-Paris, 1 dead specimen, Bouchet, Metivier and Richer de Forges collectors (Figures 10, 11). (24°45'S, 168°09'E), dredged in 230 m (127 fms), N.O. *Coriolis* CHALCAL Cruise 2, Sta. CP20, 27 October 1986, MNHN-Paris, 1 dead specimen, Bouchet, Metivier and Richer de Forges collectors.

**Remarks:** The aperture of each of the three specimens shows some wear and the shells may have been occupied by hermit crabs after the death of the snails.

**Range:** Previously known only from moderately deep water (137–154 m) off southeastern Australia (New South Wales) and the Kermadec Islands (north of New Zealand). Here recorded from the Coral Sea and off New Caledonia, in depths of 230 and 390 m (based on dead specimens).

### BIOGEOGRAPHICAL CONSIDERATIONS

The genus *Morum* (*sensu lato*) was more widely distributed during the Tertiary with species known from the Paleogene of southern Europe, India, and Java, and from the Neogene of southern Europe, Japan, New Zealand, and the middle Americas. The surviving members of the subgenera *Oniscidia* and *Herculea* are most numerous represented in the western Pacific Ocean (see range data in Table 1). Here most of the species live at moderate shelf-depths to upper slope-depths, ranging from southern Japan, the Ryukyu Archipelago, Taiwan, and the southern Philippines. In the south Pacific, a few species are known from off southeastern Australia, New Zealand, New Guinea, New Caledonia, in the Solomon, Fiji and Tonga Islands, and at Pitcairn Island, occurring in depths of 25 to 350 m. Elsewhere in the Pacific Basin, a single species of *Oniscidia* is known in Micronesia (Marshall Islands) and two occur in the tropical eastern Pacific at Cocos Island and in the Galápagos Islands. One species of this subgenus is known from the western Indian Ocean and another in the Andaman Sea. The only other members of *Oniscidia* are known from the tropical western Atlantic, where three or four species are now recognized. *Morum* (*sensu stricto*) is limited to the New World tropics, with the oldest records dating from the Plio-Pleistocene. Three species of the nominate subgenus survive, two in the Caribbean region and one in the eastern Pacific, in shallow depths.

The fossil and modern distribution of *Morum* (*sensu lato*), therefore, suggests that these gastropods are remnants of a Tethyan faunal element with somewhat limited biological dispersal potential. Although data on larval development in *Morum* are meager, these gastropods apparently do not have teleplanic larvae. *Morum* (*Morum*) *oniscus* (Linnaeus, 1767) has a non-pelagic developmental mode. In this species, the large eggs are few in number and the juveniles emerge from the capsule at the crawling stage, with a protoconch of about 2.2 mm in height and with about 1½ to 2 nuclear whorls (WORK, 1969:657, fig. 2; HUGHES & EMERSON, 1987:357, fig. 7). Living representatives of the subgenera *Oniscidia* and *Herculea*, however,










Figure 14

A living specimen of *Morum* (*O.*) *macdonaldi*, the only known "Pacific Plate" species of *Morum*, from Kwajalein Atoll, Marshall Islands; about 5 times natural size. (Drawing by Nancy Tunnell, 1988; courtesy of D. J. MacDonald.)

have larger multiwhorled protoconchs (ABBOTT, 1968:20, pl. 5). Some have 3 to 3½ nuclear whorls and a flared lip which demarcates the protoconch from the teleconch (Figures 15–20). Dr. M. G. Harasewych examined the protoconch of a specimen of *M. kurzi* (AMNH 213707; Figures 15, 16) and inferred this species to be at least lecithotrophic with a limited pelagic phase and possibly planktotrophic, on the basis of the morphology of the protoconch. He concluded: "Analysis by SHUTO's (1974) criteria [developmental mode inferred by ratio of the maximum diameter (D) to the number of protoconch whorls (VOL)] is inconclusive (VOL = 3.0; D = 1.41 mm; D/VOL = 0.47)" (Harasewych, *in litt.*, 15 May 1989). Dr. Rudolf S. Scheltema commented on the protoconchs of the three species figured herein as follows: "*Morum kurzi* (Figs. 15–16) has a planktotrophic veliger larvae as is inferred from the diameter of the apical protoconch whorl (ca. 220 μ by my measurement using the scale) and from the varix on the outer lip (which allows a rough estimate of size at settlement). The mode of development of *Morum bruuni* (Figs. 17–18) is less explicable; the rather large initial whorl (400 μ) probably indicates a direct development, presumably within the egg capsule. *Morum ponderosum* (Figs. 19–20) also must have a planktotrophic veliger. Judging by the size at settlement (estimated from the SEM micrographs) both *M. kurzi* and *M. ponderosum* probably have a planktotrophic development of at least one month (probably even 2). While 2 months is somewhat less than that of most teleplanic larvae, it nevertheless will allow considerable opportunity for passive dispersal by advection of ocean currents and may explain why *M. ponderosum* occurs on Pitcairn Island (a geologically young island of







less than 1 my) for which otherwise there is no ready explanation" (Scheltema, *in litt.*, 19 July 1989). The restricted distributional patterns indicated for other species, however, may reflect the presence of temporally limited pelagic larval stages.

The occurrences on the lithospheric plates for the 16 nominal species of *Morum* living in the Indo-West Pacific biogeographical region are tabulated (see Table 1). One of these taxa, *M. watsoni* Dance & Emerson, 1967, a new name for *M. cithara* (Watson, 1881), not *M. cythara* (Brocchi, 1814), was based on an immature specimen for which the taxonomic status is not certain. Of the remaining 15 taxa, 10 occur on the Eurasian Plate, whereas 7 or possibly 8 of these extend onto the adjacent Philippine Plate, and 3 of these also occur on the Indian-Australian Plate. Two taxa are limited, respectively, to the Indian-Australian Plate and the African Plate, and two are restricted to the Eurasian Plate. The remaining taxa are of special biogeographical interest and are discussed in more detail below.

Only 3 of the 15 recognized species (20%) are known from the Pacific Plate (Figure 1). One of these, *Morum macdonaldi*, occurs only at Kwajalein Atoll, Marshall Islands, where living specimens have been observed by SCUBA divers in 15–30 m off the exposed side of fringing reefs (Figure 14). This is the only known "low island" species. It may have been overlooked elsewhere owing to the small size of the shells (10–18 mm) and to a nocturnal habit. The same circumstances may be true for *M. kurzi*, which is well known from the southern Philippines (Philippine Plate) in tangle-net collections and is reported here from 25–30 m at Guadalcanal Island, Solomon Islands (Pacific Plate). Populations of one species, *M. ponderosum*, are apparently isolated on the Pacific Plate at Pitcairn Island, where they occur in depths to 110 m (USNM 789326). This taxon is also known from the Tonga Islands (CORNFIELD, 1986:9), New Caledonia (BOUCHET, 1981; BM[NH] 1964504), off Queensland, Australia (BEU, 1976:224) on the Indian-Australian Plate, and from the Ryukyu Archipelago (EMERSON, 1977:85) on the Philippine Plate.

As indicated above, only 1 of the 15 western Pacific species (*Morum macdonaldi*) is restricted to the Pacific Plate (Figure 1) and, thus, can be termed a "Pacific Plate species" (*cf.* SPRINGER, 1982; KAY, 1980, 1984; NEWMAN, 1986; BRIGGS, 1987). In three cases—*M. teramachii*, *M. uchiyamai*, and *M. joelgreenei*—species occur on the eastern border of the Philippine Plate in the Mariana Islands, but the deep Mariana Trench apparently prevents the non-

planktotrophic larvae of these gastropods from reaching the Pacific Plate. This is in contrast to the Mariana shoal-water faunal elements with apparent teleplanic larval development, which are believed to be a major source via a dispersal pattern for planktonically derived northwestern constituents in the Hawaiian fauna. As VERMEIJ *et al.* (1983) have noted, an offshoot of the Kuroshio Current forms a countercurrent and flows through the northern Mariana Islands, continues eastward to near Johnson Island, and eventually reaches the northwestern Hawaiian Islands (*cf.* ZINSMEISTER & EMERSON, 1979). Thus, the Mariana Trench does not seem to be a barrier for passive dispersal of hemipelagic organisms that apparently have established Pacific Plate populations via the Marianas. The expansive deep-water barrier between the Mariana Islands and the Hawaiian Archipelago does, however, serve to select against the dispersal westward of certain molluscan groups, especially archaeogastropods and intertidal species (VERMEIJ *et al.*, 1983).

The Marianas apparently originated in the mid-Paleogene to the south and perhaps west of their present position, *i.e.*, closer to the Indonesian-Philippine arc systems (KARIG, 1975). Accordingly, the distance required for dispersal of continental species to the proto-Marianas would have been reduced. Perhaps other islands (now subducted) on the Pacific Plate could have served as stepping stones for colonization by larval dispersal of the more westerly situated islands on the Pacific Plate during the Neogene (SPRINGER, 1982).

The other two species of *Morum* known from the Pacific Plate are *M. kurzi* and *M. ponderosum* (Figure 1). The former occurs in the Solomon Islands adjoining the border of the Indian-Australian Plate. It is not known from the neighboring Bismarck Archipelago and the New Hebrides Islands (Vanuatu), which also border the Indian-Australian Plate. Collections from these areas, however, are not extensive. The vast Solomon Trench may act as a barrier for dispersal into the Melanesian region for species that undergo intracapsular metamorphosis or have a brief larval stage. The presence of *M. ponderosum* in the Tonga Islands on the far western border of the Indian-Australian Plate may reflect the survival of populations that were established on the Lau-Tonga Ridge, which may have been continuous with Kermadec Island Ridge (SPRINGER, 1982). This ridge fronts a deep trench on the east face and extends southwestward to near New Zealand. This may have also served as the dispersal pathway to account for the presence of *M. bruuni* in the Kermadec Islands. *Morum cancellatum*

#### Explanations of Figures 15 to 20

Figures 15–20. Scanning electron micrographs showing lateral (left) and apical (right) views of three species of *Morum*;  $\times 50$ . Figures 15, 16: *M. (O.) kurzi*, off Davao, Mindanao Island, Philippines (AMNH 213707). Figures 17, 18: *M. (O.) bruuni*, south of New Caledonia, 24°45'S, 168°09'E (MNHN-Paris). Figures 19, 20: *M. (H.) ponderosum*, off Seragaki Beach, Okinawa (AMNH 203693).



Table 1  
Lithospheric Plate distribution of *Morum* in the Indo-West Pacific Tropics.

	n = taxa per plate				
	1 African	10 Eurasian	9 Philippine	6 Indian- Australian	3 Pacific
Genus <i>Morum</i> Roeding, 1798					
Subgenus <i>Oniscidia</i> Moersch, 1852					
1. <i>M. amabile</i> Shikama, 1973 Taiwan and ?Philippine Islands		X	X		
2. <i>M. bruuni</i> Powell, 1958 SE Australia, off New Zealand and off New Caledonia				X	
3. <i>M. cancellatum</i> G. B. Sowerby I, 1824 Southern Japan to Philippine Islands, and Fiji Islands		X	X	X	
4. <i>M. exquisitum</i> (A. Adams & Reeve, 1848) Okinawa Island and southern Philippine Islands		X	X		
5. <i>M. grande</i> (A. Adams, 1855) Southern Japan to Philippine Islands, Indonesia, and SE Aus- tralia		X	X	X	
6. <i>M. joelgreenei</i> Emerson, 1981 Southern Philippine Islands and Mariana Islands		X	X		
7. <i>M. kurzi</i> Petuch, 1979 Southern Philippine Islands and Solomon Islands			X		X
8. <i>M. macandrewi</i> G. B. Sowerby III, 1889 Southern Japan		X			
9. <i>M. macdonaldi</i> Emerson, 1981 Marshall Islands					X
10. <i>M. ninomiyai</i> Emerson, 1986 Off SW Thailand		X			
11. <i>M. praeclarum</i> Melvill, 1919 Off Somali Republic, SE Africa, Mozambique, and Seychelle Islands	X				
12. <i>M. teramachii</i> Kuroda & Habe in Habe, 1961 Southern Japan, Philippine Islands and Mariana Islands		X	X		
13. <i>M. watanabei</i> Kosuge, 1981 Southern Philippines and possibly southern Japan		X	?		
14. <i>M. watsoni</i> Dance & Emerson, 1967 Kai Islands, off New Guinea				X	
15. <i>M. uchiyamai</i> Kuroda & Habe in Habe, 1961 Southern Japan, Taiwan, Mariana Islands, and New Caledonia		X	X	X	
Subgenus <i>Herculea</i> Hanley in H. & A. Adams, 1858					
16. <i>M. ponderosum</i> (Hanley, 1858) Ryukyu Archipelago, Okinawa Island, SE Australia (Queens- land), New Caledonia, Tonga Islands, and Pitcairn Island			X	X	X

also occurs on the Indian-Australian Plate where it borders the Pacific Plate near the Fiji Islands. The apparent isolation of *M. ponderosum* on the southeastern Pacific Plate at Pitcairn Island could suggest a relict-faunal element, or more likely reflects larval recruitment from more westerly situated populations.

BOUCHET & POPPE (1988:24–30) discussed the biogeographical significance of the volutid genera *Alcithoe* and *Lyria* in the New Caledonian region, where *Morum bruuni* and *M. uchiyamai* are here recorded. As appears to be the case for some of the species of *Morum*, these volutes apparently also have non-planktotrophic larval development. Bouchet & Poppe concluded that the Norfolk Ridge, which

extends from New Zealand to New Caledonia, was a probable pathway for the dispersal of *Alcithoe*. In the case of *Lyria*, they presented a scenario for demersal dispersal, which seems unlikely at these depths (230–400 m), rather than one primarily based on vicariant events.

The genus *Morum* was recently allocated to the family Harpidae within a new subfamily, Moruminae (HUGHES & EMERSON, 1987; EMERSON & HUGHES, 1988). Both *Harpa* (Harpinae) and *Morum* have well-documented post-Tethyan fossil records, but *Harpa* may differ from *Morum* in the mode of development. The larvae of *Harpa*, in contrast to *Morum*, are believed to have an extended planktotrophic stage. In *Harpa*, the large capsules contain nu-



merous eggs (3000 to 4000 per capsule) and the protoconch consists of  $3\frac{1}{4}$  to  $5\frac{1}{4}$  whorls (HUGHES & EMERSON, 1987: 357; TAYLOR 1975:394, pl. 67; REHDER, 1973:237, pl. 208). *Harpa* is known from the Neogene of France, Italy, the Caribbean, Peru, Fiji, and Japan. Like *Morum*, most of the living species of *Harpa*, i.e., 8 of the 10 recognized species, are largely concentrated in the Indo-Pacific biogeographical region, with one species each in tropical eastern Pacific and eastern Atlantic waters (REHDER, 1973). Unlike *Morum*, however, 50% of the Indo-Pacific species extend eastward onto the Pacific Plate. Three of the four Pacific Plate *Harpa* (*H. harpa* Linnaeus, 1758; *H. amouretta* Roeding, 1798; and *H. major* Roeding, 1798) have extensive geographical ranges, occurring from east Africa to the Hawaiian Islands, with the last two taxa also known from the Marquesas Islands (REHDER, 1973; KAY, 1979; SALVAT & RIVES, 1980). The fourth of these species (*H. gracilis* Broderip & Sowerby, 1829) may be restricted to the Pacific Plate (Ellice Islands [Vuvalu] to the Tuamotus Archipelago and Marquesas Islands [AMNH 240922], and on Clipperton Island, REHDER, 1973:243), although a specimen of this species was recently attributed to Guam, in the Mariana Islands (teste Albert Deynzer, 1988). It is notable that the three species of *Harpa* (*H. harpa*, *H. amouretta*, and *H. major*) previously reported from Guam (teste B. D. Smith, 1986) are also found in the Hawaiian Islands (KAY, 1979:284). The Hawaiian fauna, however, lacks *Morum* as well as representatives of the Volutidae, Cancellariidae, Turbinellidae, and other groups of prosobranch gastropods that are mostly limited to a direct developmental mode (cf. SCHELTEMA, 1986:253).

The sharply contrasting distributional patterns of *Morum* and *Harpa* seemingly reflect the limited biological dispersal potential of most of the species of *Morum*. With the demise of the circumtropical Tethyan seaways, *Morum* was largely restricted in the Pacific tropics to the continental margins and the larger islands off the continental borderlands. This pattern of distribution has existed despite the fact that the major current circulation in the tropical Pacific has favored east to west dispersal pathways commencing in the late Neogene or perhaps much earlier (cf. NEWTON, 1988), whereas the Tethyan circulation was predominately west to east (cf. GRIGG, 1988).

#### ACKNOWLEDGMENTS

I thank the following for calling my attention to critically important specimens in their respective collections: Philippe Bouchet of the Muséum National d'Histoire Naturelle, Paris (New Caledonia); Robert B. Moffitt of the National Marine Fisheries Service Honolulu Laboratory (Mariana Islands); and Don Pisor of Pisor's Marine Shells, San Diego (Solomon Islands). The following kindly provided me with information and/or lent me specimens: Alan Beu of the New Zealand Geological Survey; Rüdiger Bieker and Betty Jean Piech of the Delaware Museum of Natural History, Wilmington; George M. Davis and M.

A. Garback of the Academy of Natural Sciences of Philadelphia; Albert Deynzer of Showcase Shells, Sanibel, FL; M. G. Harasewych and Raye N. Germon of the National Museum of Natural History, Washington, D.C.; Robert Foster of Abbey Specimen Shells, Santa Barbara; Silvard Kool, Museum of Comparative Zoology, Harvard University; James H. McLean of the Natural History Museum of Los Angeles County; and Barry D. Smith of the University of Guam Marine Laboratory. I am grateful to my AMNH colleagues, Walter E. Sage, III for technical assistance, Andrew S. Modell for photographic services, Peling Fong for SEM assistance, and Stephanie Crooms for word-processing the manuscript. I am indebted to Philippe Bouchet, Henry W. Chaney of the Santa Barbara Museum of Natural History, M. G. Harasewych, Robert B. Moffitt, and Rudolf S. Scheltema of the Woods Hole Oceanographic Institution for reviewing the manuscript and offering helpful suggestions. The conclusions, however, remain the responsibility of the author.

#### LITERATURE CITED

- ABBOTT, R. T. 1968. The helmet shells of the world. (Cassidae) Part 1. Indo-Pacific Mollusca 2(9):15-202.
- ADAMS, A. 1855. Description of new genera and species of gasteropodous [sic] Mollusca. Proc. Zool. Soc. London for 1853, 21(259):182-189 (16 May 1855).
- ADAMS, A. & L. REEVE. 1848[-50]. Mollusca. x + 87 pp. In: A. Adams, The zoology of the voyage of H.M.S. *Samarang*; under the command of Captain Sir Edward Belcher. London.
- ADAMS, H. & A. ADAMS. 1853-58. The genera of Recent Mollusca; arranged according to their organization. London. 3 Vols. 661 pp., 138 pls. (for dates of publication: see Vol. 2, p. 661).
- BEU, A. G. 1976. Revision of the southwest Pacific species of *Morum* (*Oniscidia*) (Gastropoda: Cassidae). Jour. Malacol. Soc. Australia 3(3-4):223-231.
- BOUCHET, P. 1981. Recent finds. Rossiniana, Bull. Assoc. Conchyl. Nouvelle-Calédonie, No. 10:12.
- BOUCHET, P. 1986. Oceanographic campaigns in New Caledonia. Rossiniana, Bull. Assoc. Conchyl. Nouvelle-Calédonie, No. 31:7, 8.
- BOUCHET, P. & G. T. POPPE. 1988. Deep water volutes from the New Caledonian region with a discussion on biogeography. Venus, Malacol. Soc. Japan 47(1):15-32.
- BRIGGS, J. C. 1987. Biogeography and plate tectonics. Chapter 14, The oceanic plates. Pp. 140-156. In: Development in palaeontology and stratigraphy. Elsevier: New York. 10:xi + 204 pp.
- BRODERIP, W. J. & G. B. SOWERBY. 1829. Observations on new or interesting Mollusca contained, for the most part, in the Museum of the Zoological Society. Zool. Jour., London 4:359-379.
- BRONN, H. G. 1849. Handbuch der Geschichte der Natur. 3(3) Index Palaeontologicus. Stuttgart. 980 pp.
- CORNFIELD, G. 1986. Afloat and ashore in the South Pacific. Hawaiian Shell News 33(2):9.
- DANCE, S. P. & W. K. EMERSON. 1967. Notes on *Morum dennisoni* and related species. Veliger 10(2):91-98.
- EMERSON, W. K. 1977. Notes on some Indo-Pacific species of *Morum* (Gastropoda: Tonnacea). Nautilus 91(3):81-86.



- EMERSON, W. K. 1981. Two new Indo-Pacific species of *Morum* (Gastropoda: Tonnacea). *Nautilus* 95(3):101-105.
- EMERSON, W. K. 1985. Remarks on some western Pacific species of *Morum* (Gastropoda: Tonnacea). Pp. 51-56. In: J. M. Lindsay (ed.), *Stratigraphy, palaeontology, malacology papers in honour of Dr. Nell Ludbrook*. Spec. Publ. So. Aust. Dept. Mines and Energy 5.
- EMERSON, W. K. 1986. A new species of *Morum* from the Andaman Sea (Gastropoda: Volutacea). *Nautilus* 100(3): 96-98.
- EMERSON, W. K. & R. N. HUGHES. 1988. Harpidae Bronn, 1849 (Gastropoda): conserved by ICZN. *Veliger* 30(4):423-424.
- EMERSON, W. K. & R. B. MOFFITT. 1988. A new genus and species of prosobranch gastropod (?Fascioliidae) from the Mariana Islands. *Veliger* 31(1/2):43-45.
- GARRARD, T. A. 1961. Mollusca collected by M. V. "Challenger" off the coast of Australia. *Jour. Malacol. Soc. Australia* 1(5):2-37.
- GLASS, C. & R. FOSTER. 1986. A pictorial review of the genus *Morum*. *Amer. Conchologist (Bull. Conchologists of America)* 14(4):67-69.
- GLASS, C. & R. FOSTER. 1987. Addendum to "A pictorial review of the genus *Morum*." *Amer. Conchologist (Bull. Conchologists of America)* 15(2):10.
- GRIGG, R. W. 1988. Paleooceanography of coral reefs in the Hawaiian-Emperor chain. *Science* 240(4860):1737-1743.
- HABE, T. 1961. Coloured illustrations of the shells of Japan. 2. Hoikusha, Osaka. 183 pp.
- HANLEY, S. 1858. Description of a new *Oniscia*. *Proc. Zool. Soc. London* 26(142):255-256.
- HUGHES, R. N. & W. K. EMERSON. 1987. Anatomical and taxonomic characteristics of *Harpa* and *Morum* (Neogastropoda: Harpidae). *Veliger* 29(4):349-358.
- INABA, T. & K. OYAMA. 1977. Catalogue of molluscan taxa described by Tadashige Habe during 1939-1975, with illustrations of hitherto unfigured species. Tokyo. 185 pp.
- KARIG, D. E. 1975. Basin genesis in the Philippine Sea. *Initial Reports Deep Sea Drilling Proj.* 31:857-879.
- KAY, E. A. 1979. Hawaiian marine shells. Reef and shore fauna of Hawaii. Sec. 4, Mollusca. Bernice P. Bishop Mus. Spec. Pub. 64(4):i-xviii+1-653.
- KAY, E. A. 1980. Little worlds of the Pacific: an essay on Pacific basin biogeography. Harold L. Lyon Arboretum Lecture No. 9, University of Hawaii.
- KAY, E. A. 1984. Patterns of speciation in the Indo-West Pacific. Pp. 15-31. In: P. Raven, F. Radovsky & S. H. Sohmer (eds.), *Biogeography of the tropical Pacific: proceedings of a symposium*. Assoc. System. Coll. and B. P. Bishop Museum.
- KOSUGE, S. 1981. Descriptions of two new species of the genus *Morum* with remarks on the Recent species from Philippines. *Bull. Inst. Malacol., Tokyo* 1(7):101-104.
- LINNAEUS, C. 1758. *Systema naturae per regna tria naturae. Editio decima*. Vol. 1, Regnum animale. Stockholm. 824 pp.
- LINNAEUS, C. 1767. *Systema naturae per regna tria naturae. Editio duodecima, reformata*. Vol. 1, Regnum animale. Pt. 2, pp. 533-1327 Stockholm.
- MELVILL, J. C. 1919. Descriptions of *Morum praeclarum*, sp. nov., with remarks on the Recent species of the genus. *Proc. Malacol. Soc. London* 13(3-4):69-72; Addendum, *Ibid.* 13(5-6):145.
- MOERCH, O. A. L. 1852. *Catalogus conchyliorum quae reliquit D. Alphonso D'Aguirra & Gadea, Comes Regis Daniae. Hafniae: Copenhagen. Fac. 1.* 170 pp.
- NEWMAN, W. A. 1986. Origin of the Hawaiian marine fauna: dispersal and vicariance as indicated by barnacles and other organisms. Pp. 21-49. In: R. H. Gore & K. L. Heck (eds.), *Crustacean biogeography*. A. A. Balkema: Boston.
- NEWTON, C. R. 1988. Significance of "Tethyan" fossils in the American Cordillera. *Science* 242(4877):385-391.
- PETUCH, E. J. 1979. Twelve new Indo-Pacific gastropods. *Nemouria, Occ. Pap. Delaware Mus. Natur. Hist.*, No. 23: 21 pp.
- PETUCH, E. J. 1987. New Caribbean molluscan faunas. Coastal Education and Research Foundation: Charlottesville, Virginia. 154 + 4 pp. [published 10 July 1987, *teste* editor].
- POWELL, A. W. B. 1958. Mollusca of the Kermadec Islands. Pt. I. Records Auckland Inst. Mus. 5(1-2):65-85.
- REHDER, H. A. 1973. The family Harpidae of the world. *Indo-Pacific Mollusca* 3(6):207-221.
- RICHER DE FORGES, B. 1988. Le navire oceanographique "Vauban." *Rossiniana, Bull. Assoc. Conchyl. Nouvelle-Calédonie*, No. 38:5, 6.
- ROEDING, P. F. 1798. *Museum Boltenianum sive catalogus cimeliorum e tribus regnis naturae . . .*, pt. 2, Conchyli. Hamburg. vii + 109 pp.
- SALVAT, B. & C. RIVES. 1980. *Conquillages de Polynésie. Les éditions du Pacifique*. 2nd ed. 391 pp.
- SCHELTEMA, R. S. 1986. Long-distance dispersal by planktonic larvae of shoal-water benthic invertebrates among central Pacific islands. *Bull. Mar. Sci.* 39(2):241-256.
- SHIKAMA, T. 1973. Description of new marine Gastropoda from the East and South China Seas. *Scient. Rept. Yokohama Natl. Univ.*, (Sec. 2, Biol. and Geol. Sci.) 20:1-8.
- SHUTO, T. 1974. Larval ecology of prosobranch gastropods and its bearing on biogeography and paleontology. *Lethaia* 7(3): 239-256.
- SOWERBY, G. B., I. 1824. *Genera of Recent and fossil shells. London. Vol. 1, Oniscia*, pl. 233.
- SOWERBY, G. B., III. 1889. Descriptions of fourteen new species of shells from China, Japan, and the Andaman Islands. *Proc. Zool. Soc. London for 1888*:565-570.
- SPRINGER, V. G. 1982. Pacific plate biogeography with special reference to shorefishes. *Smithsonian Contrib. Zool.* 367:1-182.
- TAYLOR, J. B. 1975. Planktonic prosobranch veligers of Kaneohe Bay. Ph.D. Dissertation, University of Hawaii. [Facsimile reprint, Univ. Microfilms International, 1982:xiii + 593.]
- VERMEIJ, G. J., E. A. ALLISON & L. G. ELDREDGE. 1983. Mollusca of the northern Mariana Islands, with special reference to the selectivity of oceanic dispersal barriers. *Micronesica* 19(1-2):27-55.
- WORK, R. C. 1969. Systematics, ecology, and distribution of the mollusks of Los Roques, Venezuela. *Bull. Mar. Sci.* 19(3):614-711.
- ZINSMEISTER, W. J. & W. K. EMERSON. 1979. The role of passive dispersal in the distribution of hemipelagic invertebrates, with examples from the tropical Pacific Ocean. *Veliger* 22(1):32-40.





Emerson, William K. 1990. "NEW RECORDS FOR WESTERN PACIFIC MORUM (GASTROPODA, HARPIDAE) WITH BIOGEOGRAPHIC IMPLICATIONS." *The veliger* 33, 145–154.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/134373>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/94220>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: California Malacozoological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.