

Range Extensions of Mollusk Species Found on the Tropical Coast of El Salvador

BY

MARCO ANTONIO HERNÁNDEZ

Curador de la Sección de Malacología y de Invertebrados Marinos del Museo de Historia Natural de El Salvador

THE COASTS OF EL SALVADOR form part of the tropical Pacific coastline of Central America, making it part of the Panamic Province. This province extends from Cabo San Lucas (Gulf of California, approximately 30°30'N) to Cabo Blanco, north of Perú (4°15'S).

It is possible that the coasts of this region are some of the most representative of the malacological fauna in the American Pacific. STUARDO (1964) estimated the diversity of this area to be 2 200 species, including the gastropods, pelecypods, scaphopods and polyplacophorans. KEEN (1971), studying the same tropical zone, found that the diversity of mollusks is much higher and reports approximately 3 317 species.

Early in 1977, the Museo de Historia Natural de El Salvador started in a systematic manner a study aimed at an inventory of the malacological fauna of the country. To date, this work has identified close to 250 species, 26 of which are not previously reported in El Salvador (Tables 1 and 2).

Generally, the species of mollusks that inhabit the coasts of El Salvador live in environments that are relatively uniform, with gradual climatological fluctuations. STUARDO (1964) suggests that among the environmental factors having the greatest influence on these invertebrates is temperature. OLSSON (1961) reports the surface temperature of the waters of the Panamic Province as varying between 26.6° and 29.4°C, with the exception of temporary cold water upwellings.

The coasts of El Salvador are characterized by their volcanic origin; some parts demonstrate the geological history of the area (GIERLOFF-EMDEN, 1971).

The sandy beaches present a topography in which strong waves remove great masses of sand, and set up surface currents; for example, Costa del Sol, El Pimiental and Los Blancos.

Of the rocky beaches, most representative are Los Cóbano and Maculís.

Table 1

Extensions of range northward to the coast of El Salvador

Species	New northern record
Gastropoda	
<i>Tegula pellisserpentis</i> (Wood)	Los Cóbano, Sonsonate
<i>Turbo saxosus</i> (Wood)	El Tamarindo, estuary
<i>Astraea buschii</i> (Philippi)	El Tamarindo, estuary
<i>Littorina varia</i> Sowerby	Barra de Santiago and El Tamarindo
<i>Cerithidea pulchra</i> (C. B. Adams)	El Tamarindo, estuary
<i>Anachis rugosa</i> (Sowerby)	Solimar; also El Tamarindo
<i>Microcithara cithara</i> (Reeve)	Los Cóbano, Sonsonate
<i>Cancellaria bulbulus</i> Sowerby	Costa del Sol; also, El Tamarindo
<i>Conus patricius</i> (Hinds)	Costa del Sol; also, El Tamarindo
Pelecypoda	
<i>Anadara similis</i> (C. B. Adams)	Los Cóbano; also, El Tamarindo
<i>Chama buddiana</i> C. B. Adams	Maculís, La Unión
<i>Protothaca beili</i> (Olsson)	El Tamarindo, estuary
<i>Tellina ecuadoriana</i> Pilsbry & Olsson	El Tamarindo, estuary
<i>Strigilla disjuncta</i> (Carpenter)	El Tamarindo, estuary
<i>Donax dentifer</i> Hanley	Barra de Santiago; also, El Tamarindo
<i>Corbula tumaca</i> (Olsson)	El Tamarindo, estuary
<i>Periploma pentadactylus</i> Pilsbry & Olsson	Costa del Sol; La Paz

Added in Proof

Gastropoda*Cyclothyca corrugata* Stearns Playa El Zonte, La Libertad

Table 2

Extensions of range southward to the coast of El Salvador

Species	New southern record
Gastropoda	
<i>Astraea olivacea</i> (Wood)	Los Cóbano; Costa del Sol; El Tamarindo
<i>Cerithium maculosum</i> (Kiener)	Los Cóbano, Sonsonate
<i>Planaxis obsoletus</i> Menke	Los Cóbano and Maculís, La Unión
<i>Acanthina tyrianthina</i> Berry	Los Cóbano, Sonsonate
<i>Solenosteira gatesi</i> Berry	Barra de Santiago and El Tamarindo
<i>Northia northiae</i> (Griffith & Pidgeon)	Costa del Sol and El Tamarindo
<i>Fusinus ambustus</i> (Gould)	Julupita, La Libertad, estuary
Pelecypoda	
<i>Cardita affinis</i> Sowerby	El Tamarindo and Maculís, La Unión
<i>Amphichaena kindermanni</i> Philippi	Metalío and El Tamarindo

Los Cóbano is characterized by having the formation of a true reef. The beach is strewn with large quantities of organic material (shell fragments mixed with sand).

In the Golfo de Fonseca (El Tamarindo, Playitas and Maculís) relatively tranquil waters provide favorable conditions for an abundant invertebrate fauna and some marine mammals.

Literature Cited

- GIERLOFF-EMDEN, H. G.
1971. La Costa de El Salvador. Prim. Edic. Direcc. Public. del Ministerio de Educ. El Salvador: 273 pp.
- KEEN, A. MYRA
1971. Sea shells of tropical West America: marine mollusks from Baja California to Peru, 2nd ed. Stanford Univ. Press, Stanford, Calif. i-xiv+1064 pp.; ca. 4000 text figs.; 22 col. pls. (21 September 1971)
- OLSSON, AXEL ADOLF
1961. Mollusks of the tropical eastern Pacific, particularly from the southern part of the Panamic-Pacific faunal province (Panama to Peru). Panamic-Pacific Pelecypoda. Paleont. Res. Inst. Ithaca, N. Y., 574 pp., 86 pls.
- STUARDO, JOSÉ
1964. Distribución de los moluscos marinos litorales en Latinoamérica. Bol. Inst. Biol. Mar. del Mar del Plata no. 7: 79-91

Added in Proof

Pelecypoda

Plicatula anomioides Keen El Pital, La Libertad



Hernandez, Ma. 1979. "RANGE EXTENSIONS OF MOLLUSK SPECIES FOUND ON THE TROPICAL COAST OF EL-SALVADOR." *The veliger* 22, 204–205.

View This Item Online: <https://www.biodiversitylibrary.org/item/134937>

Permalink: <https://www.biodiversitylibrary.org/partpdf/97664>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: California Malacozoological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.